



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>C12N 9/50, 15/57, C11D 7/42</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 91/00345</b>  <b>(43) International Publication Date:</b> 10 January 1991 (10.01.91)
<b>(21) International Application Number:</b> PCT/DK90/00164 <b>(22) International Filing Date:</b> 26 June 1990 (26.06.90)  <b>(30) Priority data:</b> 3169/89                      26 June 1989 (26.06.89)                      DK  <b>(71) Applicant:</b> NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK).  <b>(72) Inventors:</b> BRANNER, Sven ; Ved Smedebakken 7A, DK-2800 Lyngby (DK). HASTRUP, Sven ; Frederiksborgvej 10, 3th, DK-2400 København N.V (DK). OLSEN, Ole, Hvilsted ; Bækkeskovvej 38, DK-2700 Brønshøj (DK). NØRSKOV-LAURITSEN, Leif ; Birkevej 10, Strøbyegede, DK-4600 Køge (DK). SIMONSEN, Merete ; Blegdamsvej 114, 4th, DK-2100 København Ø (DK). AAS-LYNG, Dorrit ; Fyrren 8, Svogerslev, DK-4000 Roskilde (DK). CASTELEIJN, Eric ; Palletierburg 105, NL-2907 CG Cappelle a/d IJssel (NL). EGMOND, Maarten, Robert ; De Ness 34, NL-3461 GD Linschoten (NL). HAV-ERKAMP, Johan ; De Kroeskarper 40, NL-2661 KL Bergschenhoek (NL). MARUGG, John, David ; Fivelingo 141, NL-3524 BL Utrecht (NL). MOOREN, Arnoldus, Theodorus, Anthonius ; Zwaluwenlaan 461, NL-3136 TP Vlaardingen (NL).		<b>(81) Designated States:</b> AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE*, DE (European patent)*, DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> A MUTATED SUBTILISIN PROTEASE  <b>(57) Abstract</b>  Enzymes produced by mutating the genes for a number of subtilisin proteases and expressing the mutated genes in suitable hosts are presented. The mutated enzymes have a changed net electrostatic charge compared to the parent protease at the same pH. The enzymes exhibit improved wash performance in comparison to their wild type parent enzymes. The enzymes are well-suited for use in detergent compositions.		

### DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

#### *FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark			US	United States of America

**Title:****A MUTATED SUBTILISIN PROTEASE.**5 FIELD OF THE INVENTION

This invention relates to novel mutant enzymes or enzyme variants useful in formulating detergent compositions in exhibiting improved wash performance, cleaning and detergent compositions containing said enzymes, mutated genes coding for the expression of said enzymes when inserted in a suitable host cell or organism, and methods of selecting the amino acid residues to be changed in a parent enzyme in order to perform better in a given wash liquor under specified conditions.

15

BACKGROUND OF THE INVENTION

In the detergent industry enzymes have for more than 20 years been implemented in washing formulations. Enzymes used in such formulations comprise proteases, lipases, amylases, cellulases, as well as other enzymes, or mixtures thereof. Commercially most important are proteases.

Although proteases have been used in the detergent industry for more than 20 years, it is still not exactly known which physical or chemical characteristics are responsible for a good washing performance or ability of a protease.

The currently used proteases have been found by isolating proteases from nature and testing them in detergent formulations.

30

BACILLUS PROTEASES

Enzymes cleaving the amide linkages in protein substrates are classified as proteases, or (interchangeably) peptidases (see Walsh, 1979, Enzymatic Reaction Mechanisms. W.H. Freeman and Company, San Francisco, Chapter 3). Bacteria of the *Bacillus* species secrete two extracellular species of protease, a neutral, or metalloprotease, and an alkaline protease which is functionally a serine endopeptidase, referred to as subtilisin. Secretion of these proteases has been linked to the

bacterial growth cycle, with greatest expression of protease during the stationary phase, when sporulation also occurs. Joliffe et al. (1980, J. Bacterial 141:1199-1208) has suggested that *Bacillus* proteases function in cell wall turnover.

5

#### SUBTILISIN

A serine protease is an enzyme which catalyses the hydrolysis of peptide bonds, and in which there is an essential serine residue at the active site (White, Handler and  
10 Smith, 1973 "Principles of Biochemistry," Fifth Edition, McGraw-Hill Book Company, NY, pp. 271-272).

The bacterial serine proteases have molecular weights in the 20,000 to 45,000 range. They are inhibited by diisopropylfluorophosphate, but in contrast to metalloproteases, are  
15 resistant to ethylene diamino tetraacetic acid (EDTA) (although they are stabilized at high temperatures by calcium ions). They hydrolyse simple terminal esters and are similar in activity to eukaryotic chymotrypsin, also a serine protease. A more narrow term, alkaline protease, covering a  
20 sub-group, reflects the high pH optimum of some of the serine proteases, from pH 9.0 to 11.0 (for review, see Priest, 1977, Bacteriological Rev. 41:711-753).

In relation to the present invention a subtilisin is a serine protease produced by Gram-positive bacteria or fungi.  
25 A wide variety of subtilisins have been identified, and the amino acid sequence of a number of subtilisins have been determined. These include at least six subtilisins from Bacillus strains, namely, subtilisin 168, subtilisin BPN', subtilisin Carlsberg, subtilisin DY, subtilisin amylosacchariticus, and mesentericopeptidase (Kurihara et al., 1972, J.Biol.Chem. 247:5629-5631; Wells et al., 1983, Nucleic Acids Res. 11:7911-7925; Stahl and Ferrari, 1984, J.Bacteriol. 159:811-819, Jacobs et al., 1985, Nucl.Acids Res. 13:8913-8926; Nedkov et al., 1985, Biol.Chem. Hoppe-Seyler 366:421-35 430, Svendsen et al., 1986, FEBS Lett 196:228-232), one subtilisin from an actinomycetales, thermitase from Thermoactinomyces vulgaris (Meloun et al., 1985, FEBS Lett. 1983: 195-200), and one fungal subtilisin, proteinase K from Tri-

tirachium album (Jany and Mayer, 1985, Biol.Chem. Hoppe-Seyler 366:584-492).

Subtilisins are well-characterized physically and chemically. In addition to knowledge of the primary structure (amino acid sequence) of these enzymes, over 50 high resolution X-ray structures of subtilisin have been determined which delineate the binding of substrate, transition state, products, at least three different protease inhibitors, and define the structural consequences for natural variation (Kraut, 1977, Ann.Rev.Biochem. 46:331-358).

In the context of this invention, a subtilisin variant or mutated subtilisin protease means a subtilisin that has been produced by an organism which is expressing a mutant gene derived from a parent microorganism which possessed an original or parent gene and which produced a corresponding parent enzyme, the parent gene having been mutated in order to produce the mutant gene from which said mutated subtilisin protease is produced when expressed in a suitable host.

Random and site-directed mutations of the subtilisin gene have both arisen from knowledge of the physical and chemical properties of the enzyme and contributed information relating to subtilisin's catalytic activity, substrate specificity, tertiary structure, etc. (Wells et al., 1987, Proc.Natl.Acad.Sci. U.S.A. 84; 1219-1223; Wells et al., 1986, Phil.Trans.R.Soc.Lond.A. 317:415-423; Hwang and Warshel, 1987, Biochem. 26:2669-2673; Rao et al., 1987, Nature 328:551-554).

Especially site-directed mutagenesis of the subtilisin genes has attracted much attention, and various mutations are described in the following patent applications and patents:

EP publ. no. 130756 (GENENTECH) (corresponding to US Patent No. 4,760,025 (GENENCOR)) relating to site specific or randomly generated mutations in "carbonyl hydrolases" and subsequent screening of the mutated enzymes for various properties, such as  $k_{cat}/K_m$  ratio, pH-activity profile, and oxidation stability. Apart from revealing that site-specific mutation is feasible, and that mutation of subtilisin BPN' in certain specified positions, i.e. <sup>1</sup>Tyr, <sup>32</sup>Asp, <sup>155</sup>Asn, <sup>104</sup>Tyr, <sup>222</sup>Met, <sup>166</sup>Gly, <sup>64</sup>His, <sup>169</sup>Gly, <sup>189</sup>Phe, <sup>33</sup>Ser, <sup>221</sup>Ser, <sup>217</sup>Tyr, <sup>156</sup>Glu or <sup>152</sup>Ala, provide for enzymes exhibiting altered properties, this

application does not contribute to solving the problem of deciding where to introduce mutations in order to obtain enzymes with desired properties.

EP publ. no. 214435 (HENKEL) relating to cloning and expression of subtilisin Carlsberg and two mutants thereof. In this application no reason for mutation of <sup>158</sup>Asp to <sup>158</sup>Ser and <sup>161</sup>Ser to <sup>161</sup>Asp is provided.

In International patent publication No. WO 87/04461 (AMGEN) it is proposed to reduce the number of Asn-Gly sequences present in the parent enzyme in order to obtain mutated enzymes exhibiting improved pH and heat stabilities, in the application emphasis is put on removing, mutating, or modifying the <sup>109</sup>Asn and the <sup>218</sup>Asn residues in subtilisin BPN'.

International patent publication No. WO 87/05050 (GENEX) discloses random mutation and subsequent screening of a large number of mutants of subtilisin BPN' for improved properties. In the application mutations are described in positions <sup>218</sup>Asn, <sup>131</sup>Gly, <sup>254</sup>Thr, <sup>166</sup>Gly, <sup>116</sup>Ala, <sup>188</sup>Ser, <sup>126</sup>Leu, and <sup>53</sup>Ser.

In EP application no. 87303761 (GENENTECH) it is described how homology considerations at both primary and tertiary structural levels may be applied to identify equivalent amino acid residues whether conserved or not. This information together with the inventors knowledge of the tertiary structure of subtilisin BPN' lead the inventors to select a number of positions susceptible to mutation with an expectation of obtaining mutants with altered properties. The positions so identified are: <sup>124</sup>Met, <sup>222</sup>Met, <sup>104</sup>Tyr, <sup>152</sup>Ala, <sup>156</sup>Glu, <sup>166</sup>Gly, <sup>169</sup>Gly, <sup>189</sup>Phe, <sup>217</sup>Tyr. Also <sup>155</sup>Asn, <sup>21</sup>Tyr, <sup>22</sup>Thr, <sup>24</sup>Ser, <sup>32</sup>Asp, <sup>33</sup>Ser, <sup>36</sup>Asp, <sup>46</sup>Gly, <sup>48</sup>Ala, <sup>49</sup>Ser, <sup>50</sup>Met, <sup>77</sup>Asn, <sup>87</sup>Ser, <sup>94</sup>Lys, <sup>95</sup>Val, <sup>96</sup>Leu, <sup>107</sup>Ile, <sup>110</sup>Gly, <sup>170</sup>Lys, <sup>171</sup>Tyr, <sup>172</sup>Pro, <sup>197</sup>Asp, <sup>199</sup>Met, <sup>204</sup>Ser, <sup>213</sup>Lys, and <sup>221</sup>Ser, which positions are identified as being expected to influence various properties of the enzyme. Also, a number of mutations are exemplified to support these suggestions. In addition to single mutations in these positions the inventors also performed a number of multiple mutations. Further the inventors identify <sup>215</sup>Gly, <sup>67</sup>His, <sup>126</sup>Leu, <sup>135</sup>Leu, and amino acid residues within the segments 97-103, 126-129, 213-

215, and 152-172 as having interest, but mutations in any of these positions are not exemplified.

EP publ. no. 260105 (GENENCOR) describes modification of certain properties in enzymes containing a catalytic triad by selecting an amino acid residue within about 15 Å from the catalytic triad and replace the selected amino acid residue with another residue. Enzymes of the subtilisin type described in the present specification are specifically mentioned as belonging to the class of enzymes containing a catalytic triad. In subtilisins positions 222 and 217 are indicated as preferred positions for replacement.

International patent publication No. WO 88/06624 (GIST-BROCADES NV) discloses the DNA and amino acid sequences of a subtilisin protease designated PB92 almost 100% homologous in the amino acid sequence to the amino acid sequence of Subtilisin 309

International patent publication No. WO 88/07578 (GENENTECH) claims mutated enzymes derived from a precursor enzyme by replacement or modification of at least one catalytic group of an amino acid residue. The inventors state that by doing so a mutated enzyme is obtained that is reactive with substrates containing the modified or replaced catalytic group (substrate-assisted catalysis).

The general theory is based on B. amyloliquefaciens subtilisin (BPN'), where modifications have been described in positions <sup>64</sup>His that was modified into <sup>64</sup>Ala alone or in combination with a "helper" mutation of Ser-24-Cys. Modifications are also suggested in the amino acid residues <sup>32</sup>Asp, and <sup>221</sup>Ser, and a "helper" mutation of Ala-48-Glu.

International patent publication No. WO 88/08028 (GENEX) discloses genetic engineering around metal ion binding sites for the stabilization of proteins. This publication also uses Subtilisin BPN' as example and points at the following amino acid residues as candidates for substitution <sup>172</sup>Pro (P172D, P172E), <sup>131</sup>Gly (G131D), <sup>76</sup>Asn (N76D; N76D+P172D(E)), <sup>78</sup>Ser (S78D). Further, suggestions are made for the combined mutants N76D+S78D+G131D+P172D(E); N76D+G131D; S78D+G131D; S78D+P172D(E) AND S78D+G131D+P172D(E)

International patent publication No. WO 88/08033 (AMGEN) discloses a number of subtilisin analogues having a modified calcium binding site and either Asn or Gly replaced in any Asn-Gly sequence present in the molecule thereby obtaining enzymes exhibiting improved thermal and pH stability. One of the calcium binding sites is disclosed as involving the amino acid residues <sup>41</sup>Asp, <sup>75</sup>Leu, <sup>76</sup>Asn, <sup>77</sup>Asn, <sup>78</sup>Ser, <sup>79</sup>Ile, <sup>80</sup>Gly, <sup>81</sup>Val, <sup>208</sup>Thr, and <sup>214</sup>Tyr; other potential calcium binding sites are suggested at <sup>140</sup>Asp, and <sup>172</sup>Pro; <sup>14</sup>Pro, and <sup>271</sup>Gln; and <sup>172</sup>Pro and <sup>195</sup>Glu or <sup>197</sup>Asp. Also mutating the <sup>109</sup>Asn and <sup>218</sup>Asn positions is suggested. Mutants produced are N109S, N109S+N218S, N76D+N109S+N218S, N76D+N77D+N109S+N218S, N76D+I79E+N109S+N218S.

International patent publication No. WO 88/08164 (GENEX) describes a method for identifying residues in a protein which may be substituted by a cysteine to permit formation of potentially stabilizing disulfide bonds. The method is based on detailed knowledge of the three dimensional structure of the protein and uses a computer for selecting the positions. In relation to subtilisin proteases, Subtilisin BPN' was used as a model system. Using the method on Subtilisin BPN' resulted in the suggestion of 11 candidates for introducing disulfide bonds (1:T22C+S87C, 2:V26C+L235C, 3:G47C+P57C, 4:M50C+N109C, 5:E156C+T164C, 6:V165C+K170C, 7:V165C+S191C, 8:Q206C+A216C, 9:A230C+V270C, 10:I234C+A274C, 11:H238C+W241C). Of these 4 were produced (1, 2, 4, and 8) of which 2 did not provide any stabilizing effect (2 and 4). Further mutants were produced by combining two of these mutants with each other, and one with another mutation, viz. T22C+S87C+N218S, and T22C+S87C+Q206C+A216C. Also, a number of further unsuccessful mutants were produced, viz. A1C+S78C, S24C+S87C, K27C+S89C, A85C+A232C, I122C+V147C, S249C+A273C, and T253C+A273C.

Also, it has been shown by Thomas, Russell, and Fersht, Nature 318, 375-376(1985) that exchange of <sup>99</sup>Asp into <sup>99</sup>Ser in subtilisin BPN' changes the pH dependency of the enzyme.

In a subsequent article J. Mol. Biol. (1987) 193, 803-813, the same authors also discuss the substitution of <sup>156</sup>Ser in place of <sup>156</sup>Glu.

Both these mutations are within a distance of about 15Å from the active <sup>64</sup>His.



In Nature 328, 496-500(1987) Russel and Fersht discuss the results of their experiments and present rules for changing pH-activity profiles by mutating an enzyme to obtain changes in surface charge.

5

#### ISOELECTRIC POINT ( $pI_0$ )

The isoelectric point,  $pI_0$ , is defined as the pH value where the enzyme molecule complex (with optionally attached metal or other ions) is neutral, i. e. the sum of electrostatic charges (net electrostatic charge, =NEC) on the complex is equal to zero. In this sum of course consideration of the positive or negative nature of the individual electrostatic charges must be taken into account.

The isoelectric point is conveniently calculated by using equilibrium considerations using pK values for the various charged residues in the enzyme in question and then by iteration find the pH value where the NEC of the enzyme molecule is equal to zero.

One problem with this calculation is that the pK values for the charged residues are dependent on their environment and consequently subject to variation. However, very good results are obtainable by allocating specific approximate pK values to the charged residues independently of the actual value. It is also possible to perform more sophisticated calculations, partly taking the environment into consideration.

The  $pI_0$  may also be determined experimentally by isoelectric focusing or by titrating a solution containing the enzyme. Also, the various pK values for the charged residues may be determined experimentally by titration.

30

#### INDUSTRIAL APPLICATIONS OF SUBTILISINS

Proteases such as subtilisins have found much utility in industry, particularly in detergent formulations, as they are useful for removing proteinaceous stains.

At present the following proteases are known and many of them are marketed in large quantities in many countries of the world.

Subtilisin BPN<sup>1</sup> or Novo, available from e.g. SIGMA, St. Louis, U.S.A.

8

Subtilisin Carlsberg, marketed by Novo-Nordisk a/s (Denmark) as ALCALASE® and by IBIS (Holland) as MAXATASE®;

A Bacillus lentus subtilisin, marketed by NOVO INDUSTRI A/S (Denmark) as SAVINASE®;

5 Enzymes closely resembling SAVINASE® such as MAXACAL® marketed by IBIS, and OPTICLEAN® marketed by MILES KALI CHEMIE (FRG);

A Bacillus lentus subtilisin, marketed by Novo-Nordisk a/s (Denmark) as ESPERASE®;

10 KAZUSASE® marketed by SHOWA DENKO (Japan)

To be effective, however, such enzymes must not only exhibit activity under washing conditions, but must also be compatible with other detergent components during detergent production and storage.

15 For example, subtilisins may be used in combination with other enzymes active against other substrates, and the selected subtilisin should possess stability towards such enzymes, and also the selected subtilisin preferably should not catalyse degradation of the other enzymes. Also, the chosen  
20 subtilisin should be resistant to the action from other components in the detergent formulation, such as bleaching agents, oxidizing agents, etc., in particular an enzyme to be used in a detergent formulation should be stable with respect to the oxidizing power, calcium binding properties, and pH  
25 conditions rendered by the non-enzymatic components in the detergent during storage and in the wash liquor during wash.

The ability of an enzyme to catalyse the degradation of various naturally occurring substrates present on the objects to be cleaned during e.g. wash is often referred to as  
30 its washing ability, washability, detergency, or wash performance. Throughout this application the term wash performance will be used to encompass this property.

Naturally occurring subtilisins have been found to  
35 possess properties which are highly variable in relation to their washing power or ability under variations in parameters such as pH. Several of the above marketed detergent proteases, indeed, have a better performance than those marketed about 20 years ago, but for optimal performance each enzyme has its own

specific conditions regarding formulation and wash conditions, e.g. pH, temperature, ionic strength (=I), active system (tensides, surfactants, bleaching agent, etc.), builders, etc.

As a consequence it is found that an enzyme possessing desirable properties at low pH and low I may be less attractive at more alkaline conditions and high I, or an enzyme exhibiting fine properties at high pH and high I may be less attractive at low pH, low I conditions.

The advent and development of recombinant DNA techniques has had a profound influence in the field of protein chemistry.

It has been envisaged that these techniques will make it possible to design peptides and proteins, such as enzymes, and hormones according to desired specifications, enabling the production of compounds exhibiting desired properties.

It is possible now to construct enzymes having desired amino acid sequences, and as indicated above a fair amount of research has been devoted to designing subtilisins with altered properties. Among the proposals the technique of producing and screening a large number of mutated enzymes as described in EP publ. no. 130756 (GENENTECH) (US Patent No. 4,760,025 (GENENCOR)) and International patent publ. no. WO 87/05050 (GENEX) correspond to the classical method of isolating native enzymes and screening them for their properties, but is more efficient through the knowledge of the presence of a large number of different mutant enzymes.

Since a subtilisin enzyme typically comprises 275 amino acid residues each capable of being 1 out of 20 possible naturally occurring amino acids, one very serious draw-back in that procedure is the very large number of mutations generated that has to be submitted to a preliminary screening prior to further testing of selected mutants showing interesting characteristics at the first screening, since no guidance is indicated in determining which amino acid residues to change in order to obtain a desired enzyme with improved properties for the use in question, such as, in this case formulating detergent compositions exhibiting improved wash performance under specified conditions of the wash liquor.

A procedure as outlined in these patent applications will consequently only be slightly better than the traditional random mutation procedures which have been known for years.

The other known techniques relate to changing specific properties, such as transesterification and hydrolysis rate (EP publ. no. 260105 (GENENCOR)), pH-activity profile (Thomas, Russell, and Fersht, supra), and substrate specificity (International patent publ. no. WO 88/07578 (GENENTECH)). None of these publications relates to changing the wash performance of enzymes.

A further technique that has evolved is using the detailed information on the three dimensional structure of proteins for analyzing the potential consequences of substituting certain amino acids. This approach has been used and is described in EP 260105 (GENENCOR), WO 88/07578 (GENENTECH), WO 88/08028 (GENEX), WO 88/08033 (AMGEN), and WO 88/08164 (GENEX)

Thus, as indicated above no relationship has yet been identified between well defined properties of an enzyme such as those mentioned above and the wash performance of an enzyme.

In unpublished International Patent Application no. PCT/DK 88/00002 (NOVO INDUSTRI A/S) it is proposed to use the concept of homology comparison to determine which amino acid positions should be selected for mutation and which amino acids should be substituted in these positions in order to obtain a desired change in wash performance.

By using such a procedure the task of screening is reduced drastically, since the number of mutants generated is much smaller, but with that procedure it is only foreseen that enzymes exhibiting the combined useful properties of the parent enzyme and the enzyme used in the comparison may be obtained.

The problem seems to be that although much research has been directed at revealing the mechanism of enzyme activity, still only little is known about the factors in structure and amino acid residue combination that determine the properties of enzymes in relation to their wash performance.

Consequently there still exists a need for further improvement and tailoring of enzymes to wash systems, as well

as a better understanding of the mechanism of protease action in the practical use of cleaning or detergent compositions. Such an understanding could result in rules which may be applied for selecting mutations that with a reasonable degree of certainty will result in an enzyme exhibiting improved wash performance under specified conditions in a wash liquor.

#### SUMMARY OF THE INVENTION

Further investigations into these problems have now surprisingly shown that one of the critical factors in the use of subtilisin enzymes in detergent compositions is the adsorption of the enzyme to the substrate, i.e. the material to be removed from textiles, hard surfaces or other materials to be cleaned.

Consequently, the present invention relates to mutations of the subtilisin gene resulting in changed properties of the mutant subtilisin enzyme expressed by such a mutated gene, whereby said mutant subtilisin enzyme exhibit improved behaviour in detergent compositions. Mutations are generated at specific nucleic acids in the parent subtilisin gene responsible for the expression of specific amino acids in specific positions in the subtilisin enzyme.

The present invention also relates to methods of selecting the positions and amino acids to be mutated, and thereby mutatis mutandis the nucleic acids to be changed in the subtilisin gene in question.

The invention relates, in part, but is not limited to, mutations of the subtilisin 309 and subtilisin Carlsberg genes and ensuing mutant subtilisin 309 and Carlsberg enzymes, which exhibit improved wash performance in different detergent compositions resulting in wash liquors of varying pH values.

Further the invention relates to the use of the mutant enzymes in cleaning compositions and cleaning compositions comprising the mutant enzymes, especially detergent compositions comprising the mutant subtilisin enzymes.

ABBREVIATIONSAMINO ACIDS

	A =	Ala	=	Alanine
	V =	Val	=	Valine
5	L =	Leu	=	Leucine
	I =	Ile	=	Isoleucine
	P =	Pro	=	Proline
	F =	Phe	=	Phenylalanine
	W =	Trp	=	Tryptophan
10	M =	Met	=	Methionine
	G =	Gly	=	Glycine
	S =	Ser	=	Serine
	T =	Thr	=	Threonine
	C =	Cys	=	Cysteine
15	Y =	Tyr	=	Tyrosine
	N =	Asn	=	Asparagine
	Q =	Gln	=	Glutamine
	D =	Asp	=	Aspartic Acid
	E =	Glu	=	Glutamic Acid
20	K =	Lys	=	Lysine
	R =	Arg	=	Arginine
	H =	His	=	Histidine

NUCLEIC ACID BASES

25	A =	Adenine	
	G =	Guanine	
	C =	Cytosine	
	T =	Thymine	(only in DNA)
	U =	Uracil	(only in RNA)

30

MUTATIONS

In describing the various mutants produced or contemplated according to the invention, the following nomenclatures were adapted for ease of reference:

35 Original amino acid(s) position(s) substituted amino acid(s)

According to this the substitution of Glutamic acid for glycine in position 195 is designated as:

Gly 195 Glu or G195E

a deletion of glycine in the same position is:

13

Gly 195 \* or G195\*

and insertion of an additional amino acid residue such as lysine is:

Gly 195 GlyLys or G195GK

5 Where a deletion is indicated in Table I, or present in a subtilisin not indicated in Table I, an insertion in such a position is indicated as:

\* 36 Asp or \*36D

for insertion of an aspartic acid in position 36

10 Multiple mutations are separated by pluses, i.e.:

Arg 170 Tyr + Gly 195 Glu or R170Y+G195E

representing mutations in positions 170 and 195 substituting tyrosine and glutamic acid for arginine and glycine, respectively.

14  
TABLE ICOMPARISON OF AMINO ACID SEQUENCE FOR VARIOUS PROTEASES

5

- a = subtilisin BPN' (Wells et al, 1983, supra)  
b = subtilisin amylosacchariticus (Kurihara et al, 1972, supra)  
10 c = subtilisin 168 (Stahl and Ferrari, 1984, supra)  
d = subtilisin mesentericopeptidase (Svendsen et al, 1986, supra)  
e = subtilisin DY (Nedkov et al, 1985, supra)  
f = subtilisin Carlsberg (Smith et al, 1968, supra)  
15 g = subtilisin Carlsberg (Jacobs et al, 1985, supra)  
h = subtilisin 309 (International Patent Application No. PCT/DK 88/00002)  
i = subtilisin 147 (International Patent Application No. PCT/DK 88/00002)  
20 j = thermitase (Meloun et al, 1985, supra)  
k = proteinase K (Betz et al, 1988, Eur. J. Biochem. 178: 155 ff), and Gunkel et al, 1989, Eur. J. Biochem. 179: 185 ff)  
l = aqualysin (Kwon et al, 1988, Eur. J. Biochem. 173:491 ff)  
25 m = Bacillus PB92 protease (European Patent Publication No. 0 283 075)  
n = Protease TW7 (Tritirachium album) (International Patent Application No. PCT/US88/01040)  
30 o = Protease TW3 (Tritirachium album) (International Patent Application No. PCT/US88/01040)  
\* = assigned deletion

35

continued....



....Table I continued

No:	1	10	
5	a) *-*-*-*-A-Q-S--V-P-Y-G-V-S-Q-I-K-***-A-P-A-		
	b) *-*-*-*-A-Q-S--V-P-Y-G-I-S-Q-I-K-***-A-P-A-		
	c) *-*-*-*-A-Q-S--V-P-Y-G-I-S-Q-I-K-***-A-P-A-		
	d) *-*-*-*-A-Q-S--V-P-Y-G-I-S-Q-I-K-***-A-P-A-		
	e) *-*-*-*-A-Q-T--V-P-Y-G-I-P-L-I-K-***-A-D-K-		
10	f) *-*-*-*-A-Q-T--V-P-Y-G-I-P-L-I-K-***-A-D-K-		
	g) *-*-*-*-A-Q-T--V-P-Y-G-I-P-L-I-K-***-A-D-K-		
	h) *-*-*-*-A-Q-S--V-P-W-G-I-S-R-V-Q-***-A-P-A-		
	i) *-*-*-*-Q-T--V-P-W-G-I-S-F-I-N-***-T-Q-Q-		
	j) Y-T-P-N-D-P-Y-F-S-S--R-Q-Y-G-P-Q-K-I-Q-***-A-P-Q-		
15	k) *-*-*-*-A-A-Q-T-N-A-P-W-G-L-A-R-I-S-S-T-S-P-G-T-S-T-		
	l) *-*-*-*-A-T-Q-S-P-A-P-W-G-L-D-R-I-D-Q-R-D-L-P-L-S-N-		
	m) *-*-*-*-A-Q-S--V-P-W-G-I-S-R-V-Q-***-A-P-A-		
	n) *-*-*-*-A-T-Q-E-D-A-P-W-G-L-A-R-I-S-S-Q-E-P-G-G-T-T-		
	o) *-*-*-*-A-E-Q-R-N-A-P-W-G-L-A-R-I-S-S-T-S-P-G-T-S-T-		
20			
No:	20	30	40
	a) L-H-S-Q-G-Y-T-G-S-N-V-K-V-A-V-I-D-S-G-I-D-S-S-H-P-D-L-*		
	b) L-H-S-Q-G-Y-T-G-S-N-V-K-V-A-V-I-D-S-G-I-D-S-S-H-P-D-L-*		
	c) L-H-S-Q-G-Y-T-G-S-N-V-K-V-A-V-I-D-S-G-I-D-S-S-H-P-D-L-*		
25	d) L-H-S-Q-G-Y-T-G-S-N-V-K-V-A-V-I-D-S-G-I-D-S-S-H-P-D-L-*		
	e) V-Q-A-Q-G-Y-K-G-A-N-V-K-V-G-I-I-D-T-G-I-A-A-S-H-T-D-L-*		
	f) V-Q-A-Q-G-F-K-G-A-N-V-K-V-A-V-L-D-T-G-I-Q-A-S-H-P-D-L-*		
	g) V-Q-A-Q-G-F-K-G-A-N-V-K-V-A-V-L-D-T-G-I-Q-A-S-H-P-D-L-*		
	h) A-H-N-R-G-L-T-G-S-G-V-K-V-A-V-L-D-T-G-I-*S-T-H-P-D-L-*		
30	i) A-H-N-R-G-I-F-G-N-G-A-R-V-A-V-L-D-T-G-I-*A-S-H-P-D-L-*		
	j) A-W-*D-I-A-E-G-S-G-A-K-I-A-I-V-D-T-G-V-Q-S-N-H-P-D-L-A-		
	k) Y-Y-Y-D-E-S-A-G-Q-G-S-C-V-Y-V-I-D-T-G-I-E-A-S-H-P-E-F-*		
	l) S-Y-T-Y-T-A-T-G-R-G-V-N-V-Y-V-I-D-T-G-I-R-T-T-H-R-E-F-*		
	m) A-H-N-R-G-L-T-G-S-G-V-K-V-A-V-L-D-T-G-I-*S-T-H-P-D-L-*		
35	n) Y-T-Y-D-D-S-A-G-T-G-T-C-A-Y-I-I-D-T-G-I-Y-T-N-H-T-D-F-*		
	o) Y-R-Y-D-D-S-A-G-Q-G-T-C-V-Y-V-I-D-T-G-V-E-A-S-H-P-E-F-*		
			continued...

continued....

....Table I continued

No:	50	60	
5	a) *-K-V-A-G-G-A-S-M-V-P-S-E-T-N-P-F-*-Q-D-N-N-S-H-G-T-H-V-		
	b) *-N-V-R-G-G-A-S-F-V-P-S-E-T-N-P-Y-*-Q-D-G-S-S-H-G-T-H-V-		
	c) *-N-V-R-G-G-A-S-F-V-P-S-E-T-N-P-Y-*-Q-D-G-S-S-H-G-T-H-V-		
	d) *-N-V-R-G-G-A-S-F-V-P-S-E-T-N-P-Y-*-Q-D-G-S-S-H-G-T-H-V-		
	e) *-K-V-V-G-G-A-S-F-V-S-G-E-S-*-Y-N-*-T-D-G-N-G-H-G-T-H-V-		
10	f) *-N-V-V-G-G-A-S-F-V-A-G-E-A-*-Y-N-*-T-D-G-N-G-H-G-T-H-V-		
	g) *-N-V-V-G-G-A-S-F-V-A-G-E-A-*-Y-N-*-T-D-G-N-G-H-G-T-H-V-		
	h) *-N-I-R-G-G-A-S-F-V-P-G-E-P-*-S-T-*-Q-D-G-N-G-H-G-T-H-V-		
	i) *-R-I-A-G-G-A-S-F-I-S-S-E-P-*-S-Y-*-H-D-N-N-G-H-G-T-H-V-		
	j) G-K-V-V-G-G-W-D-F-V-D-N-D-S-T-P-*-Q-N-G-N-G-H-G-T-H-C-		
15	k) *-E-G-R-A-Q-M-V-K-T-Y-Y-Y-S-S-*-R-D-G-N-G-H-G-T-H-C-		
	l) *-G-G-R-A-R-V-G-Y-D-A-L-G-G-N-G-*-Q-D-C-N-G-H-G-T-H-V-		
	m) *-N-I-R-G-G-A-S-F-V-P-G-E-P-*-S-T-*-Q-D-G-N-G-H-G-T-H-V-		
	n) *-G-G-R-A-K-F-L-K-N-F-A-G-D-G-Q-D-T-D-G-N-G-H-G-T-H-V-		
	o) *-E-G-R-A-Q-M-V-K-T-Y-Y-A-S-S-*-R-D-G-N-G-H-G-T-H-C-		
20			
No:	70	80	90
	a) A-G-T-V-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-S-A-S-L-Y-A-V-K-V-		
	b) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-S-A-S-L-Y-A-V-K-V-		
	c) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-S-P-S-A-S-L-Y-A-V-K-V-		
25	d) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-S-A-S-L-Y-A-V-K-V-		
	e) A-G-T-V-A-A-L-*-D-N-T-T-G-V-L-G-V-A-P-N-V-S-L-Y-A-I-K-V-		
	f) A-G-T-V-A-A-L-*-D-N-T-T-G-V-L-G-V-A-P-S-V-S-L-Y-A-V-K-V-		
	g) A-G-T-V-A-A-L-*-D-N-T-T-G-V-L-G-V-A-P-S-V-S-L-Y-A-V-K-V-		
	h) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-S-A-E-L-Y-A-V-K-V-		
30	i) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-S-A-D-L-Y-A-V-K-V-		
	j) A-G-I-A-A-A-V-T-N-N-S-T-G-I-A-G-T-A-P-K-A-S-I-L-A-V-R-V-		
	k) A-G-T-V-G-S-*-R-*-T-Y-G-V-A-K-K-T-Q-L-F-G-V-K-V-		
	l) A-G-T-I-G-G-V-*-T-Y-G-V-A-K-A-V-N-L-Y-A-V-R-V-		
	m) A-G-T-I-A-A-L-*-N-N-S-I-G-V-L-G-V-A-P-N-A-E-L-Y-A-V-K-V-		
35	n) A-G-T-V-G-G-T-*-T-Y-G-V-A-K-K-T-S-L-F-A-V-K-V-		
	o) A-G-T-I-G-S-*-R-*-T-Y-G-V-A-K-K-T-Q-I-F-G-V-K-V-		

continued....

....Table I continued

No:	100	110	120
5	a) L-G-A-D-G-S-G-Q-Y-S-W-I-I-N-G-I-E-W--A-I-A--N-N-M-D--		
	b) L-D-S-T-G-S-G-Q-Y-S-W-I-I-N-G-I-E-W--A-I-A--N-N-M-D--		
	c) L-D-S-T-G-S-G-Q-Y-S-W-I-I-N-G-I-E-W--A-I-A--N-N-M-D--		
	d) L-D-S-T-G-S-G-Q-Y-S-W-I-I-N-G-I-E-W--A-I-A--N-N-M-D--		
	e) L-N-S-S-G-S-G-T-Y-S-A-I-V-S-G-I-E-W--A-T-Q--N-G-L-D--		
10	f) L-N-S-S-G-S-G-S-Y-S-G-I-V-S-G-I-E-W--A-T-T--N-G-M-D--		
	g) L-N-S-S-G-S-G-T-Y-S-G-I-V-S-G-I-E-W--A-T-T--N-G-M-D--		
	h) L-G-A-S-G-S-G-S-V-S-S-I-A-Q-G-L-E-W--A-G-N--N-G-M-H--		
	i) L-D-R-N-G-S-G-S-L-A-S-V-A-Q-G-I-E-W--A-I-N--N-N-M-H--		
	j) L-D-N-S-G-S-G-T-W-T-A-V-A-N-G-I-T-Y--A-A-D--Q-G-A-K--		
15	k) L-D-D-N-G-S-G-Q-Y-S-T-I-I-A-G-M-D-F-V-A-S-D-K-N-N-R-N-C-		
	l) L-D-C-N-G-S-G-S-T-S-G-V-I-A-G-V-D-W-V--T--R-N-H-R-R-P-		
	m) L-G-A-S-G-S-G-S-V-S-S-I-A-Q-G-L-E-W--A-G-N--N-G-M-H--		
	n) L-D-A-N-G-Q-G-S-N-S-G-V-I-A-G-M-D-F-V-T-K-D-A-S-S-Q-N-C-		
	o) L-N-D-Q-G-S-G-Q-Y-S-T-I-I-S-G-M-D-F-V-A-N-D-Y-R-N-R-N-C-		
20			
No:	130	140	
	a) *-***-V-I-N-M-S-L-G-G-P-S-G-S-A-A-L-K-A-A-V-D-K-A-V-A-		
	b) *-***-V-I-N-M-S-L-G-G-P-S-G-S-T-A-L-K-T-V-V-D-K-A-V-S-		
	c) *-***-V-I-N-M-S-L-G-G-P-T-G-S-T-A-L-K-T-V-V-D-K-A-V-S-		
25	d) *-***-V-I-N-M-S-L-G-G-P-T-G-S-T-A-L-K-T-V-V-D-K-A-V-S-		
	e) *-***-V-I-N-M-S-L-G-G-P-S-G-S-T-A-L-K-Q-A-V-D-K-A-Y-A-		
	f) *-***-V-I-N-M-S-L-G-G-A-S-G-S-T-A-M-K-Q-A-V-D-N-A-Y-A-		
	g) *-***-V-I-N-M-S-L-G-G-P-S-G-S-T-A-M-K-Q-A-V-D-N-A-Y-A-		
	h) *-***-V-A-N-L-S-L-G-S-P-S-P-S-A-T-L-E-Q-A-V-N-S-A-T-S-		
30	i) *-***-I-I-N-M-S-L-G-S-T-S-G-S-S-T-L-E-L-A-V-N-R-A-N-N-		
	j) *-***-V-I-S-L-S-L-G-G-T-V-G-N-S-G-L-Q-Q-A-V-N-Y-A-W-N-		
	k) P-K-G-V-V-A-S-L-S-L-G-G-G-Y-S-S-S-V-N-S-A-A-A--R-L-Q-S-		
	l) A-***-V-A-N-M-S-L-G-G-G-V--S-T-A-L-D-N-A-V-K-N-S-I-A-		
	m) *-***-V-A-N-L-S-L-G-S-P-S-P-S-A-T-L-E-Q-A-V-N-S-A-T-S-		
35	n) P-K-G-V-V-V-N-M-S-L-G-G-P-S-S-S-A-V-N-R-A-A-A--E-I-T-S-		
	o) P-N-G-V-V-A-S-M-S-I-G-G-G-Y-S-S-S-V-N-S-A-A-A--N-L-Q-Q-		

continued....

....Table I continued

No:	150	160	170
5 a)	S-G-V-V-V-V-A-A-A-G-N-E-G-T-S-G-S-S-S-T-V-G-Y-P-G-K-Y-P-		
b)	S-G-I-V-V-A-A-A-A-G-N-E-G-S-S-G-S-S-S-T-V-G-Y-P-A-K-Y-P-		
c)	S-G-I-V-V-A-A-A-A-G-N-E-G-S-S-G-S-T-S-T-V-G-Y-P-A-K-Y-P-		
d)	S-G-I-V-V-A-A-A-A-G-N-E-G-S-S-G-S-T-S-T-V-G-Y-P-A-K-Y-P-		
e)	S-G-I-V-V-V-A-A-A-G-N-S-G-S-S-G-S-Q-N-T-I-G-Y-P-A-K-Y-D-		
10 f)	R-G-V-V-V-V-A-A-A-G-N-S-G-N-S-G-S-T-N-T-I-G-Y-P-A-K-Y-D-		
g)	R-G-V-V-V-V-A-A-A-G-N-S-G-S-S-G-N-T-N-T-I-G-Y-P-A-K-Y-D-		
h)	R-G-V-L-V-V-A-A-S-G-N-S-G-A*-G-S-I-S*-*-Y-P-A-R-Y-A-		
i)	A-G-I-L-L-V-G-A-A-G-N-T-G-R*-Q-G-V-N*-*-Y-P-A-R-Y-S-		
j)	K-G-S-V-V-V-A-A-A-G-N-A-G-N-T-A-P-N*-*-Y-P-A-Y-Y-S-		
15 k)	S-G-V-M-V-A-V-A-A-G-N-N-N-A-D-A-R-N-Y-S*-*-P-A-S-E-P-		
l)	A-G-V-V-Y-A-V-A-A-G-N-D-N-A-N-A-C-N-Y-S*-*-P-A-R-V-A-		
m)	R-G-V-L-V-V-A-A-S-G-N-S-G-A*-G-S-I-S*-*-Y-P-A-R-Y-A-		
n)	A-G-L-F-L-A-V-A-A-G-N-E-A-T-D-A-S-S-S-S*-*-P-A-S-E-E-		
o)	S-G-V-M-V-A-V-A-A-G-N-N-N-A-D-A-R-N-Y-S*-*-P-A-S-E-S-		
20			
No:	180	190	200
a)	S-V-I-A-V-G-A-V-D-S-S-N-Q-R-A-S-F-S-S-V-G-P-E-L-D-V-M-A-		
b)	S-T-I-A-V-G-A-V-N-S-S-N-Q-R-A-S-F-S-S-A-G-S-E-L-D-V-M-A-		
c)	S-T-I-A-V-G-A-V-N-S-S-N-Q-R-A-S-F-S-S-A-G-S-E-L-D-V-M-A-		
25 d)	S-T-I-A-V-G-A-V-N-S-A-N-Q-R-A-S-F-S-S-A-G-S-E-L-D-V-M-A-		
e)	S-V-I-A-V-G-A-V-D-S-N-K-N-R-A-S-F-S-S-V-G-A-E-L-E-V-M-A-		
f)	S-V-I-A-V-G-A-V-D-S-N-S-N-R-A-S-F-S-S-V-G-A-E-L-E-V-M-A-		
g)	S-V-I-A-V-G-A-V-D-S-N-S-N-R-A-S-F-S-S-V-G-A-E-L-E-V-M-A-		
h)	N-A-M-A-V-G-A-T-D-Q-N-N-N-R-A-S-F-S-Q-Y-G-A-G-L-D-I-V-A-		
30 i)	G-V-M-A-V-A-A-V-D-Q-N-G-Q-R-A-S-F-S-T-Y-G-P-E-I-E-I-S-A-		
j)	N-A-I-A-V-A-S-T-D-Q-N-D-N-K-S-S-F-S-T-Y-G-S-V-V-D-V-A-A-		
k)	S-V-C-T-V-G-A-S-D-R-Y-D-R-R-S-S-F-S-N-Y-G-S-V-L-D-I-F-G-		
l)	E-A-L-T-V-G-A-T-T-S-S-D-A-R-A-S-F-S-N-Y-G-S-C-V-D-L-F-A-		
m)	N-A-M-A-V-G-A-T-D-Q-N-N-N-R-A-S-F-S-Q-Y-G-A-G-L-D-I-V-A-		
35 n)	S-A-C-T-V-G-A-T-D-K-T-D-T-L-A-E-Y-S-N-F-G-S-V-V-D-L-L-A-		
o)	S-I-C-T-V-G-A-T-D-R-Y-D-R-R-S-S-F-S-N-Y-G-S-V-L-D-I-F-A-		

continued....

....Table I continued

No:	210	220	
5	a) P-G-V-S-I-Q-S-T-L-P-G-N-*K-*Y-G-A-Y-N-G-T-S-M-A-S-P-H-		
	b) P-G-V-S-I-Q-S-T-L-P-G-G-*T-*Y-G-A-Y-N-G-T-S-M-A-T-P-H-		
	c) P-G-V-S-I-Q-S-T-L-P-G-G-*T-*Y-G-A-Y-N-G-T-S-M-A-T-P-H-		
	d) P-G-V-S-I-Q-S-T-L-P-G-G-*T-*Y-G-A-Y-N-G-T-S-M-A-T-P-H-		
	e) P-G-V-S-V-Y-S-T-Y-P-S-N-*T-*Y-T-S-L-N-G-T-S-M-A-S-P-H-		
10	f) P-G-A-G-V-Y-S-T-Y-P-T-N-*T-*Y-A-T-L-N-G-T-S-M-A-S-P-H-		
	g) P-G-A-G-V-Y-S-T-Y-P-T-S-*T-*Y-A-T-L-N-G-T-S-M-A-S-P-H-		
	h) P-G-V-N-V-Q-S-T-Y-P-G-S-*T-*Y-A-S-L-N-G-T-S-M-A-T-P-H-		
	i) P-G-V-N-V-N-S-T-Y-T-G-N-*R-*Y-V-S-L-S-G-T-S-M-A-T-P-H-		
	j) P-G-S-W-I-Y-S-T-Y-P-T-S-*T-*Y-A-S-L-S-G-T-S-M-A-T-P-H-		
15	k) P-G-T-S-I-L-S-T-W-I-G-G-*S-*T-R-S-I-S-G-T-S-M-A-T-P-H-		
	l) P-G-A-S-I-P-S-A-W-Y-T-S-D-T-A-T-Q-T-L-N-G-T-S-M-A-T-P-H-		
	m) P-G-V-N-V-Q-S-T-Y-P-G-S-*T-*Y-A-S-L-N-G-T-S-M-A-T-P-H-		
	n) P-G-T-D-I-K-S-T-W-N-D-G-R-T-K-I-I-S-*-*G-T-S-M-A-S-P-H-		
	o) P-G-T-D-I-L-S-T-W-I-G-G-S-T-R-S-I-S-*-*G-T-S-M-A-T-P-H-		
20			
No:	230	240	250
	a) V-A-G-A-A-A-L-I-L-S-K-H-P-N-W-T-N-T-Q-V-R-S-S-L-E-N-T-T-		
	b) V-A-G-A-A-A-L-I-L-S-K-H-P-T-W-T-N-A-Q-V-R-D-R-L-E-S-T-A-		
	c) V-A-G-A-A-A-L-I-L-S-K-H-P-T-W-T-N-A-Q-V-R-D-R-L-E-S-T-A-		
25	d) V-A-G-A-A-A-L-I-L-S-K-H-P-T-W-T-N-A-Q-V-R-D-R-L-E-S-T-A-		
	e) V-A-G-A-A-A-L-I-L-S-K-Y-P-T-L-S-A-S-Q-V-R-N-R-L-S-S-T-A-		
	f) V-A-G-A-A-A-L-I-L-S-K-H-P-N-L-S-A-S-Q-V-R-N-R-L-S-S-T-A-		
	g) V-A-G-A-A-A-L-I-L-S-K-H-P-N-L-S-A-S-Q-V-R-N-R-L-S-S-T-A-		
	h) V-A-G-A-A-A-L-V-K-Q-K-N-P-S-W-S-N-V-Q-I-R-N-H-L-K-N-T-A-		
30	i) V-A-G-V-A-A-L-V-K-S-R-Y-P-S-Y-T-N-N-Q-I-R-Q-R-I-N-Q-T-A-		
	j) V-A-G-V-A-G-L-L-A-S-Q-G-R-S-*-*A-S-N-I-R-A-A-I-E-N-T-A-		
	k) V-A-G-L-A-A-Y-L-M-T-L-G-K-T-T-A-A-S-A-C-R-*Y-I-A-D-T-A-		
	l) V-A-G-V-A-A-L-Y-L-E-Q-N-P-S-A-T-P-A-S-V-A-S-A-I-L-N-G-A-		
	m) V-A-G-A-A-A-L-V-K-Q-K-N-P-S-W-S-N-V-Q-I-R-N-H-L-K-N-T-A-		
35	n) V-A-G-L-G-A-Y-F-L-G-L-G-Q-K-V-Q-G-L-*C-D-*Y-M-V-E-K-G-		
	o) V-A-G-L-A-A-Y-L-M-T-L-G-R-A-T-A-S-N-A-C-R-*Y-I-A-Q-T-A-		

continued....

continued....

....Table I continued

No:	260	270	275
5 a)	T-K-L-G-D-S-F-Y-Y--G-K-G-L-I-N-V-Q-A-A-A-Q		
b)	T-Y-L-G-D-S-F-Y-Y--G-K-G-L-I-N-V-Q-A-A-A-Q		
c)	T-Y-L-G-N-S-F-Y-Y--G-K-G-L-I-N-V-Q-A-A-A-Q		
d)	T-Y-L-G-S-S-F-Y-Y--G-K-G-L-I-N-V-Q-A-A-A-Q		
e)	T-N-L-G-D-S-F-Y-Y--G-K-G-L-I-N-V-E-A-A-A-Q		
10 f)	T-Y-L-G-S-S-F-Y-Y--G-K-G-L-I-N-V-E-A-A-A-Q		
g)	T-Y-L-G-S-S-F-Y-Y--G-K-G-L-I-N-V-E-A-A-A-Q		
h)	T-S-L-G-S-T-N-L-Y--G-S-G-L-V-N-A-E-A-A-T-R		
i)	T-Y-L-G-S-P-S-L-Y--G-N-G-L-V-H-A-G-R-A-T-Q		
j)	D-K-I-S-G-T-G-T-Y-W-A-K-G-R-V-N-A-Y-K-A-V-Q-Y		
15 k)	N-K-G-D-L-S-N-I-P-F-G-T-V-N-L-L-A-Y-N-N-Y-Q-A		
l)	T-T-G-R-L-S-G-I-G-S-G-S-P-N-R-L-L-Y-S-L-L-S-S-G-S-G		
m)	T-S-L-G-S-T-N-L-Y--G-S-G-L-V-N-A-E-A-A-T-R		
n)	L-K-D-V-I-Q-S-V-P-S-D-T-A-N-V-L-I-N-N-G-E-G-S-A		
o)	N-Q-G-D-L-S-N-I-S-F-G-T-V-N-L-A-Y-N-N-Y-Q-G		

20

#### BRIEF DESCRIPTION OF THE DRAWING

The invention is described further in detail in the following parts of this specification with reference to the drawings wherein:

Figure 1 shows the construction of plasmid pSX88,  
Figure 2 shows a restriction map of plasmid pSX88,  
Figure 3 exemplifies the construction of the mutant subtilisin 309 genes for expressing the enzymes of the invention,

Figure 4 shows the restriction map for plasmid pSX92,  
and

Figure 5 graphically demonstrates the relationship between pH of maximum performance and calculated pI of the mutant enzymes of the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

Above it was stated that the invention relates to mutated subtilisins in which the amino acid sequence has been changed through mutating the gene of the subtilisin enzyme, which it is desired to modify, in codons responsible for the expression of amino acids located on or close to the surface of the resulting enzyme.

In the context of this invention a subtilisin is defined as a serine protease produced by gram-positive bacteria or fungi. In a more narrow sense, applicable to many embodiments of the invention, subtilisin also means a serine protease of gram-positive bacteria. According to another definition, a subtilisin is a serine protease, wherein the relative order of the amino acid residues in the catalytic triad is Asp - His - Ser (positions 32, 64, and 221). In a still more specific sense, many of the embodiments of the invention relate to serine proteases of gram-positive bacteria which can be brought into substantially unambiguous homology in their primary structure, with the subtilisins listed in Table I above.

Using the numbering system originating from the amino acid sequence of subtilisin BPN', which is shown in Table I above aligned with the amino acid sequence of a number of other known subtilisins, it is possible to indicate the position of an amino acid residue in a subtilisin enzyme unambiguously. Positions prior to amino acid residue number 1 in subtilisin BPN' are assigned a negative number, such as -6 for the N-terminal Y in thermitase, or 0, such as for the N-terminal A in proteinase K. Amino acid residues which are insertions in relation to subtilisin BPN' are numbered by the addition of letters in alphabetical order to the preceding subtilisin BPN' number, such as 12a, 12b, 12c, 12d, 12e for the "insert" S-T-S-P-G in proteinase K between <sup>12</sup>Ser and <sup>13</sup>Thr.

Using the above numbering system the positions of interest are:

1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24, 25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89, 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129,

130, 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, .  
155, 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167,  
170, 171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189,  
191, 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213,  
5 214, 215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241,  
242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255,  
256, 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

#### ISOELECTRIC POINT ( $pI_o$ )

10 Assuming that the substrate under washing conditions has an electrostatic charge opposite to that of the enzyme, it might be expected that the adsorption and thus the wash performance of the enzyme to the substrate would be improved by increasing the net electrostatic charge, NEC, of the enzyme.

15 However, it was surprisingly found that a decrease in the NEC of the enzyme under such circumstances could result in an improved wash performance of the enzyme.

Stated differently, it was found that changing the isoelectric point,  $pI_o$ , of the enzyme in a direction to  
20 approach a lower pH, also shifted the pH of optimum wash performance of the enzyme to a lower value, meaning that in order to design an enzyme to a wash liquor of low pH, in which the enzyme is to be active, improvement in the wash performance of a known subtilisin enzyme may be obtained by mutating the  
25 gene for the known subtilisin enzyme to obtain a mutant enzyme having a lower  $pI_o$ .

This finding led to experiments showing that the opposite also is feasible. Meaning that a known subtilisin enzyme may also be designed for use in high pH detergents by  
30 shifting its  $pI_o$  to higher values, thereby shifting the wash performance pH optimum for the enzyme to higher pH values.

The present invention therefore in one aspect relates to mutated subtilisin proteases, wherein the net electrostatic charge has been changed in comparison to the parent protease  
35 at the same pH, and in which proteases there are, relative to said parent protease, either fewer or more positively-charged amino acid residue(s) and/or more or fewer negatively-charged amino acid residue(s), or more or fewer positively-charged amino acid residue(s) and/or fewer or more negatively-charged



amino acid residue(s) among the amino acid residues at any one or more of positions

1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24, 25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89, 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130, 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155, 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170, 171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191, 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256, 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275, and whereby said subtilisin protease has an isoelectric pH ( $pI_0$ ) lower or higher, respectively, than that of said parent protease.

In a preferred embodiment the invention relates to mutant subtilisin proteases, wherein the NEC has been changed in comparison to the parent protease at the same pH, and in which proteases there are, relative to said parent protease, either fewer or more positively-charged amino acid residue(s) and/or more or fewer negatively-charged amino acid residue(s), or either more or fewer positively-charged amino acid residue(s) and/or fewer or more negatively-charged amino acid residue(s), among the amino acid residues at any one or more of positions

1, 2, 3, 4, 14, 15, 17, 18, 20, 27, 40, 41, 43, 44, 45, 46, 51, 52, 60, 61, 62, 75, 76, 78, 79, 91, 94, 97, 100, 105, 106, 108, 112, 113, 117, 118, 129, 130, 133, 134, 136, 137, 141, 143, 144, 145, 146, 165, 173, 181, 183, 184, 185, 191, 192, 206, 209, 210, 211, 212, 216, 239, 240, 242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 255, 256, 257, 259, 263, 269, 271, 272,

and whereby said subtilisin protease has an isoelectric pH ( $pI_0$ ) lower or higher, respectively, than that of said parent protease.

In another preferred embodiment the invention relates to mutant subtilisin proteases, wherein the NEC has been changed in comparison to the parent protease at the same pH, and in which proteases there are, relative to said parent  
5 protease, either fewer or more positively-charged amino acid residue(s) and/or more or fewer negatively-charged amino acid residue(s), or either more or fewer positively-charged amino acid residue(s) and/or fewer or more negatively-charged amino acid residue(s), among the amino acid residues at any one or  
10 more of positions

1, 2, 3, 4, 14, 15, 17, 18, 20, 27, 40, 41, 43, 44, 45, 46, 51, 52, 60, 61, 62, 75, 76, 78, 79, 91, 94, 97, 100, 105, 106, 108, 112, 113, 117, 118, 129, 130, 133, 134, 136, 137, 141, 143, 144, 145, 146, 165, 173, 181, 183, 184, 185, 191, 192,  
15 206, 209, 210, 211, 212, 216, 239, 240, 242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 255, 256, 257, 259, 263, 269, 271, 272,

and at least one further mutation affecting an amino acid residue occupying a position chosen from the group of posi-  
20 tions

1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24, 25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89, 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107,  
25 108, 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130, 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155, 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170, 171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191, 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213,  
30 214, 215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256, 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

and whereby said subtilisin protease has an isoelectric pH  
35 ( $pI_0$ ) lower or higher, respectively, than that of said parent protease.

In these aspects the invention in short relates to mutant proteases in which the  $pI_0$  of the mutant protease is lower than the  $pI_0$  of the parent protease, and in which the pH

for optimum wash performance is also lower than the pH optimum for the parent protease; or mutant proteases wherein the  $pI_0$  of the mutant protease is higher than the  $pI_0$  of the parent protease, and in which the pH for optimum wash performance is also higher than the pH optimum for the parent protease.

It is generally believed (Thomas, Russell, and Fersht, supra) that kinetic properties can be influenced by changes in the electrostatic surface charge in the vicinity of the active site in the enzyme, but it has now surprisingly been found that changes in the kinetic properties of an enzyme can also be brought about by modifying surface charges remote from the active site.

Consequently the invention is also considered to embrace mutant subtilisin enzymes, wherein one or more amino acid residues in a distance of more than 15Å from the catalytic triad of said enzyme has been changed in comparison to the amino acid sequence of its parent enzyme, and in a way to provide for a mutant protease having an isoelectric point ( $=pI_0$ ) shifted in the same direction as it is desired to shift the pH for optimum wash performance of the enzyme, which pH optimum should be as close as possible to the pH of the wash liquor, wherein said mutant protease is intended for use.

According to several embodiments of the invention, there are provided mutant proteases and detergent compositions containing them, wherein the amino-acid sequence of the mutant protease contains an insertion mutation at position 36, for example to insert a negatively-charged amino-acid residue (e.g. D or E) or a neutral polar residue (such as A, Q, or N) or a positive amino-acid residue, such as R or K.

This is particularly applicable for example to subtilisin proteases 309 and 147 and PB92, and any other sequence for which the homology indicates that it naturally has an amino-acid residue missing at position 36 relative to the sequence of subtilisin BPN'.

Insertion mutants at position 36 can have further mutations, for example at one or more of positions 120, 170, 195, 235, and/or 251, and/or at position 76.

Suitable mutations at position 76 are e.g. negatively charged residues such as N76D or N76E.

5 Mutations at position 36 (especially insertion of negative or polar neutral residue) and at position 76 (substitution by negatively-charged residue) can often have stabilising effect on the mutant protease, and can be used in combination. Mutations at for example one or more of positions  
10 120, 170, 195, 235 and 251 have been found to be associated with increased enzyme activity. Even in cases where these latter mutations are associated individually with some loss of stability it can be acceptable and useful to combine them with mutations at one or both of positions 36 and 76.

15

Among useful examples of such protease mutants are those having the following mutations:

- S-021) \*36D
- 20 S-022) \*36D+R170Y+G195E+K251E
- S-023) \*36D+H120D+R170Y+G195E+K235L
- S-024) \*36D+H120D+R170Y+G195E+K235L+K251E
- S-025) \*36D+H120D+G195E+K235L
- S-235) \*36D+N76D
- 25 S-035) \*36D+N76D+H120D+G195E+K235L

Under some conditions it can be advantageous to arrange a further mutation in a mutant protease having an insertion of  
30 a negative amino-acid residue at position 36, to insert a positive charge elsewhere in the molecule. This can increase the water solubility of the resulting mutant protease, e.g. where the further mutation provides a positive charge, e.g. a positively-charged residue at position 213, e.g. T213K.

35

According to the invention it is further preferred that the mutant subtilisin enzyme represents a mutation of a parent enzyme selected from subtilisin BPN', subtilisin amylosacchariticus, subtilisin 168, subtilisin mesentericopeptidase,

subtilisin Carlsberg, subtilisin DY, subtilisin 309, subtilisin 147, thermitase, Bacillus PB92 protease, and proteinase K, preferably subtilisin 309, subtilisin 147, subtilisin Carlsberg, aqualysin, Bacillus PB92 protease, Protease TW7, or  
 5 Protease TW3.

Further preferred embodiments comprise subtilisin enzymes containing one or more of the mutations:

R10F, R10L, R10F+R45A+E89S+E136Q+R145A+D181N+R186P+E271Q,  
 10 R10F+R19Q+E89S+E136Q+R145A+D181N+E271Q+R275Q, Q12K, Q12R,  
 Q12K+P14D+T22K+N43R+Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12R+P14D+T22R+N43R+Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12K+P14D+T22K+T38K+N43R+Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R,  
 15 S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12R+P14D+T22R+T38R+N43R+Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12K+P14D+T22K+T38K+N43R+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R,  
 20 Q12R+P14D+T22R+T38R+N43R+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, P14D, P14K, P14K+\*36D, P14K+N218D, P14K+P129D, A15K, A15R, R19Q, T22K, T22R, K27R, K27V, D32\*, \*36D, \*36D+R170Y+G195E+K251E, \*36D+H120D+R170Y+G195E+K235L, \*36D+H120D+R170Y+G195E+K235L+K251E, \*36D+H120D+G195E+K235L, T38K, T38R, D41E, N43R, N43K, R45A, E53R, E53K, E53G+K235L, E54G, E54Y, Q59E, Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, D60N, N76D, E89S, E89S+K251N, Y91F, K94R, G97D, G97D+H120K, A98K, A98R, S99D, S99D+N140K, E112T,  
 30 H120K, H120D, H120D+K235L, H120D+G195E+K235L, H120D+R170Y+G195E+K235L, H120D+R170Y+G195E+K235L+K251E, P129D, E136Q, E136K, E136R, E136Q+R10L, N140D, N140K, N140R, S141K, S141R, R145A, S156E, S156E+A158R+A172D+N173K, S156E+A158R+A172D+N173K+T213R, S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, A158R, A158K, Y167V, R170Y, R170Y+G195E, R170Y+K251E, R170Y+G195E+K251E, R170Y+G195E+K235L, Y171E, Y171T, A172D, N173K, D181N, N184K, N184R, N185D, R186P, Y192V, Y192V,A, G195E, G195D, G195E+T213R, G195E+K251E, G195E+K235L, D197N, D197K, D197E, Q206D, Q206E, Y209L, T213R, T213K, Y214T,

28

Y214S, N218D, N218S, K235L, K235R, K237R, W241Y, L, W241Y+H249R,  
 W241L+H249R, N248D, H249R, K251R, K251E, K251N, T255E, S256R,  
 S256K, S259L, S259D, Y263W, S265K, S265R, E271Q, E271G, E271G+  
 K27V, E271Q, G, A272R, A272R, R275Q,  
 5 D14K, D14K+D120K, D14K+D120K+D140K, D14K+D120K+D140K+D172K,  
 K27D, K27D+D120K, E54T, E54Y, N97D, N97D+S98D, N97D+T213D,  
 S98D, S98D+T213D, D120K, D140K, S156E, D172K, T213D, N218D.

Further specific preferred embodiments are mutated  
 10 subtilisin proteases comprising one or more of the mutations:

	S001)	G195E
	S002)	G195D
	S003)	R170Y
	S004)	R170Y+G195E
15	S005)	K251E
	S006)	H120D
	S008)	H120D+G195E
	S009)	T71D
	S010)	T71D+G195E
20	S011)	R170Y+K251E
	S012)	R170Y+G195E+K251E
	S013)	T71D+R170Y+K251E
	S014)	T71D+R170Y+G195E+K251E
	S015)	K235L
25	S016)	H120D+K235L
	S017)	H120D+G195E+K235L
	S018)	G195E+K251E
	S019)	H120D+R170Y+G195E+K235L
	S020)	H120D+R170Y+G195E+K235L+K251E
30	S021)	*36D
	S022)	*36D+R170Y+G195E+K251E
	S023)	*36D+H120D+R170Y+G195E+K235L
	S024)	*36D+H120D+R170Y+G195E+K235L+K251E
	S025)	*36D+H120D+G195E+K235L
35	S026)	E136R
	S027)	E89S
	S028)	D181N
	S029)	E89S+E136R
	S030)	E89S+D181N

	S031)	D197N+E271Q
	S032)	D197N
	S033)	E271Q
	S035)	*36D+N76D+H120D+G195E+K235L
5	S041)	G195F
	S201)	N76D
	S202)	N76D+G195E
	S203)	N76D+R170Y+G195E
	S204)	H120D+G195E+K235L+K251E
10	S223)	Q59E+N76D+A98R+S99D+T213K+K235L+N248D+T255E+S256K+S259D+A272R
	S224)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+K235L+N248D+T255E+S256K+S259D+A272R
	S225)	*36D+Q59E+N76D+A98R+S99D+R170Y+S156E+A158R+A172D+N173R+K235L+N248D+T255E+S256K+S259D+ A272R
15	S226)	*36Q
	S227)	*36D+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+R170Y+G195E+K235L+N248D+T255E+S256K+S259D+ A272R
	S228)	*36D+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
20	S229)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
	S230)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
	S231)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
	S232)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
	S233)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R
	S234)	Q206D
25	S235)	*36D+N76D
	S242)	*36Q+N76D+H120D+G195E+K235L
	C001)	D14K
	C002)	D120K
30	C003)	D140K
	C004)	D14K+D120K
	C005)	K27D
	C006)	K27D+D120K
	C008)	D172K
35	C009)	D14K+D120K+D140K
	C010)	D14K+D120K+D140K+D172K
	C013)	N97D
	C014)	S98D
	C015)	T213D

	C017)	S156E
	C018)	N97D+S98D
	C019)	N97D+T213D
	C022)	S98D+T213D
5	C028)	N218D
	C100)	V51D
	C101)	E54T
	C102)	E54Y

10           In a further aspect of the invention the above observations about the  $pI_0$  are further utilized in a method for determining or selecting the position(s) and the amino acid(s) to be deleted, substituted or inserted for the amino acid(s) in a parent enzyme, whereby the selection is performed in a way  
15 whereby the calculated net electrostatic charge (=NEC) in a resulting mutant enzyme has been changed in comparison to the NEC in the parent enzyme of choice calculated at the same pH value.

20           Another way of expressing this principle covered by the invention is that the position(s) and the amino acid(s) to be deleted, substituted or inserted for the amino acid(s) in said parent enzyme is selected in a way whereby the total number of charges or total charge content (=TCC), and/or the  
25 NEC in a resulting mutant enzyme is changed in a way to provide for a mutant protease having an isoelectric point ( $=pI_0$ ) shifted in the same direction as it is desired to shift the pH for optimum wash performance of the enzyme, which pH optimum should be as close as possible to the pH of the wash liquor,  
30 wherein said mutant protease is intended for use.

          As indicated above the  $pI_0$  of a macromolecule such as an enzyme is calculated as the pH where the NEC of the molecule is equal to zero. The procedure is exemplified in the examples  
35 below, but the principles are described in more detail here.

          pK values are assigned to each potentially charged amino acid residue. Then the ratio of the occurrence of an amino acid residue at a given pH in charged or uncharged form



31

(charged/uncharged, C/U(i)) is calculated for both negative and positive charge, by using the formulae Ia and Ib:

$$C/U(i) = \exp(\ln_{10}(pH - pK_i)) \quad (\text{negative charge}) \quad (Ia)$$

5

$$C/U(i) = \exp(\ln_{10}(pK_i - pH)) \quad (\text{positive charge}) \quad (Ib)$$

respectively.

From formulae Ia and Ib it is seen that at pH equal to 10  $pK_i$ , C/U(i) is equal to 1.

Then the relative charge,  $Q_r(i)$ , or charge contribution allocated to each charged residue is calculated by using the formulae IIa and IIb:

$$15 \quad Q_r(i) = C/U(i)/(1+C/U(i)) \quad (\text{negative charge}) \quad (IIa)$$

$$Q_r(i) = -C/U(i)/(1+C/U(i)) \quad (\text{positive charge}) \quad (IIb)$$

The pH value where the sum of all the charge contributions from the charged residues is equal to zero is found by iteration or through interpolation in a sufficiently dense pH-charge sum table.

#### DETERGENT COMPOSITIONS COMPRISING THE MUTANT ENZYMES

25 The present invention also comprises the use of the mutant enzymes of the invention in cleaning and detergent compositions and such composition comprising the mutant subtilisin enzymes.

Such compositions comprise in addition to any one or 30 more of the mutant subtilisin enzymes in accordance to any of the preceding aspects of the invention alone or in combination any of the usual components included in such compositions which are well-known to the person skilled in the art.

Such components comprise builders, such as phosphate 35 or zeolite builders, surfactants, such anionic, cationic or non-ionic surfactants, polymers, such as acrylic or equivalent polymers, bleach systems, such as perborate- or amino-containing bleach precursors or activators, structurants, such as

silicate structurants, alkali or acid to adjust pH, humectants, and or neutral inorganic salts.

In several useful embodiments the detergent compositions can be formulated as follows:

a) A detergent composition formulated as a detergent powder containing phosphate builder, anionic surfactant, nonionic surfactant, acrylic or equivalent polymer, perborate bleach precursor, amino-containing bleach activator, silicate or other structurant, alkali to adjust to desired pH in use, and neutral inorganic salt.

b) A detergent composition formulated as a detergent powder containing zeolite builder, anionic surfactant, nonionic surfactant, acrylic or equivalent polymer, perborate bleach precursor, amino-containing bleach activator, silicate or other structurant, alkali to adjust to desired pH in use, and neutral inorganic salt.

c) A detergent composition formulated as an aqueous detergent liquid comprising anionic surfactant, nonionic surfactant, humectant, organic acid, caustic alkali, with a pH adjusted to a value between 9 and 10.

d) A detergent composition formulated as a nonaqueous detergent liquid comprising a liquid nonionic surfactant consisting essentially of linear alkoxyated primary alcohol, triacetin, sodium triphosphate, caustic alkali, perborate monohydrate bleach precursor, and tertiary amine bleach activator, with a pH adjusted to a value between about 9 and 10.

e) A detergent composition formulated as a detergent powder in the form of a granulate having a bulk density of at least 550 g/l, e.g. at least 600 g/l, containing anionic and nonionic surfactants, e.g. anionic surfactant and a mixture of nonionic surfactants with respective alkoxylation degrees about 7 and about 3, low or substantially zero neutral inorganic

salt, phosphate builder, perborate bleach precursor, tertiary amine bleach activator, sodium silicate, and minors and moisture.

5 f) A detergent composition formulated as a detergent powder in the form of a granulate having a bulk density of at least 600 g/l, containing anionic surfactant and a mixture of nonionic surfactants with respective alkoxylation degrees about 7 and about 3, low or substantially zero neutral inorganic  
10 salt, zeolite builder, perborate bleach precursor, tertiary amine bleach activator, sodium silicate, and minors and moisture.

g) A detergent composition formulated as a detergent  
15 powder containing anionic surfactant, nonionic surfactant, acrylic polymer, fatty acid soap, sodium carbonate, sodium sulphate, clay particles with or without amines, perborate bleach precursor, tertiary amine bleach activator, sodium silicate, and minors and moisture.

20

h) A detergent composition formulated as a detergent (soap) bar containing soap based on pan-saponified mixture of tallow and coconut oil, neutralised with orthophosphoric acid, mixed with protease, also mixed with sodium formate, borax,  
25 propylene glycol and sodium sulphate, and then plodded on a soap production line.

j) An enzymatic detergent composition formulated to give a wash liquor pH of 9 or less when used at a rate corresponding  
30 to 0.4-0.8 g/l surfactant.

k) An enzymatic detergent composition formulated to give a wash liquor pH of 8.5 or more when used at a rate corresponding to 0.4-0.8 g/l surfactant.

35

l) An enzymatic detergent composition formulated to give a wash liquor ionic strength of 0.03 or less, e.g. 0.02 or less, when used at a rate corresponding to 0.4-0.8 g/l surfactant.

m) An enzymatic detergent composition formulated to give a wash liquor ionic strength of 0.01 or more, e.g. 0.02 or more, when used at a rate corresponding to 0.4-0.8 g/l  
5 surfactant.

DETERGENT COMPOSITIONS COMPRISING MUTANT ENZYMES IN COMBINATION  
WITH LIPASE

10 It has surprisingly been found that a decrease in the isoelectric point, pI, and hence net charge of a subtilisin type protease under washing conditions, can result in not only an improved wash performance of the enzyme but also an improved compatibility with lipase.

15 It has also been surprisingly found that compatibility of protease with lipase is influenced not only by the pI but also by the positions at which the charges are located relative to the active site of the protease: The introduction of negative charge or removal of positive charge closer to the  
20 active site gives stronger improvement of compatibility of protease with lipase.

Accordingly, certain embodiments of the invention provide enzymatic detergent compositions, comprising lipase and  
25 also comprising mutated subtilisin protease, wherein the net molecular electrostatic charge of the mutated protease has been changed by insertion, deletion or substitution of amino-acid residues in comparison to the parent protease, and wherein, in said protease, there are, relative to said parent protease,  
30 fewer positively-charged amino-acid residue(s) and/or more negatively-charged amino-acid residue(s), whereby said subtilisin protease has an isoelectric pH ( $pI_0$ ) lower than that of said parent protease.

35 One preferred class of lipases for such use originates in Gram-negative bacteria, and includes e.g. lipase enzymes of the groups defined in EP 0 205 208 and 0 206 390 (both to Unilever), (hereby incorporated by reference), including

lipases immunologically related to those from certain Ps. fluorescens, P. gladioli and Chromobacter strains.

Preferred embodiments of mutant subtilisin protease enzyme for use in conjunction with lipase as aforesaid possess one or more mutations at the site of an amino-acid residue located within the range of about 15A-20A from the active site, especially for example at positions 170, 120, or 195.

The lipase can usefully be added in the form of a granular composition, (alternatively a solution or a slurry), of lipolytic enzyme with carrier material (e.g. as in EP 258068 (Novo Nordisk A/S) and Savinase® and Lipolase® products of Novo Nordisk A/S).

The added amount of lipase can be chosen within wide limits, for example 50 to 30,000 LU/g per gram of the surfactant system or of the detergent composition, e.g. often at least 100 LU/g, very usefully at least 500 LU/g, sometimes preferably above 1000, above 2000 LU/g or above 4000 LU/g or more, thus very often within the range 50-4000 LU/g and possibly within the range 200-1000 LU/g. In this specification lipase units are defined as they are in EP 258068.

The lipolytic enzyme can be chosen from among a wide range of lipases: in particular the lipases described in for example the following patent specifications, EP 214761 (Novo Nordisk A/S), EP 0 258 068 and especially lipases showing immunological cross-reactivity with antisera raised against lipase from Thermomyces lanuginosus ATCC 22070, EP 0 205 208 and EP 0 206 390 and especially lipases showing immunological cross-reactivity with antisera raised against lipase from Chromobacter viscosum var lipolyticum NRRL B-3673, or against lipase from Alcaligenes PL-679, ATCC 31371 and FERM-P 3783, also the lipases described in specifications WO 87/00859 (Gist-Brocades) and EP 0 204 284 (Sapporo Breweries). Suitable in particular are for example the following commercially available lipase preparations: Novo Lipolase®, Amano lipases CE, P, B, AP, M-AP, AML, and CES, and Meito lipases MY-30, OF, and PL, also Esterase® MM, Lipozym®, SP225, SP285, Saiken lipase, Enzeco lipase, Toyo Jozo lipase and Diosynth lipase (Trade Marks).

Genetic engineering of the enzymes can be achieved by extraction of an appropriate lipase gene, e.g. the gene for lipase from *Thermomyces lanuginosus* or from a mutant thereof, and introduction and expression of the gene or derivative thereof in a suitable producer organism such as an *Aspergillus*. The techniques described in WO 88/02775 (Novo Nordisk A/S), EP 0 243 338 (Labofina), EP 0 268 452 (Genencor) and notably EP 0 305 216 (Novo Nordisk A/S) or EP 0 283 075 (Gist-Brocades) may be applied and adapted.

Similar considerations apply *mutatis mutandis* in the case of other enzymes, which may also be present. Without limitation: Amylase can for example be used when present in an amount in the range about 1 to about 100 MU (maltose units) per gram of detergent composition, (or 0.014-1.4, e.g. 0.07-0.7, KNU/g (Novo units)). Cellulase can for example be used when present in an amount in the range about 0.3 to about 35 CEVU units per gram of the detergent composition.

The detergent compositions may furthermore include the following usual detergent ingredients in the usual amounts. They may be built or unbuilt, and may be of the zero-P type (i.e. not containing any phosphorus-containing builders). Thus the composition may contain in aggregate for example from 1-50%, e.g. at least about 5% and often up to about 35-40% by weight, of one or more organic and/or inorganic builders. Typical examples of such builders include those already mentioned above, and more broadly include alkali metal ortho, pyro, and tripolyphosphates, alkali metal carbonates, either alone or in admixture with calcite, alkali metal citrates, alkali metal nitrilotriacetates, carboxymethyloxysuccinates, zeolites, polyacetalcarboxylates and so on.

Furthermore, the detergent compositions may contain from 1-35% of a bleaching agent or a bleach precursor or a system comprising bleaching agent and/or precursor with activator therefor. Further optional ingredients are lather boosters, foam depressors, anti-corrosion agents, soil-suspending agents, sequestering agents, anti-soil redeposition

agents, perfumes, dyes, stabilising agents for the enzymes and so on.

The compositions can be used for the washing of textile materials, especially but without limitation cotton and polyester-based textiles and mixtures thereof. Especially suitable are for example washing processes carried out at temperatures of about 60-65 deg C or lower, e.g. about 30°C-35°C or lower. It can be very suitable to use the compositions at a rate sufficient to provide about e.g. 0.4-0.8 g/l surfactant in the wash liquor, although it is of course possible to use lesser or greater concentrations if desired. Without limitation it can for example be stated that a use-rate from about 3 g/l and up to about 6 g/l of the detergent formulation is suitable for use in the case when the formulations are as in the Examples.

#### METHOD FOR PRODUCING MUTATIONS IN SUBTILISIN GENES

Many methods for introducing mutations into genes are well known in the art. After a brief discussion of cloning subtilisin genes, methods for generating mutations in both random sites, and specific sites, within the subtilisin gene will be discussed.

#### CLONING A SUBTILISIN GENE

The gene encoding subtilisin may be cloned from any Gram-positive bacteria or fungus by various methods, well known in the art. First a genomic, and/or cDNA library of DNA must be constructed using chromosomal DNA or messenger RNA from the organism that produces the subtilisin to be studied. Then, if the amino-acid sequence of the subtilisin is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify subtilisin-encoding clones from a genomic library of bacterial DNA, or from a fungal cDNA library. Alternatively, a labelled oligonucleotide probe containing sequences homologous to subtilisin from another strain of bacteria or fungus could be used as a probe to identify subtilisin-encoding clones, using hybridization and washing conditions of lower stringency.

Yet another method for identifying subtilisin-producing clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming protease-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for subtilisin, such as skim milk. Those bacteria containing subtilisin-bearing plasmid will produce colonies surrounded by a halo of clear agar, due to digestion of the skim milk by excreted subtilisin.

10

#### GENERATION OF RANDOM MUTATIONS IN THE SUBTILISIN GENE

Once the subtilisin gene has been cloned into a suitable vector, such as a plasmid, several methods can be used to introduce random mutations into the gene.

15 One method would be to incorporate the cloned subtilisin gene, as part of a retrievable vector, into a mutator strain of Eschericia coli.

Another method would involve generating a single stranded form of the subtilisin gene, and then annealing the fragment of DNA containing the subtilisin gene with another DNA fragment such that a portion of the subtilisin gene remained single stranded. This discrete, single stranded region could then be exposed to any of a number of mutagenizing agents, including, but not limited to, sodium bisulfite, hydroxylamine, nitrous acid, formic acid, or hydralazine. A specific example of this method for generating random mutations is described by Shortle and Nathans (1978, Proc. Natl. Acad. Sci. U.S.A., 75: 2170-2174). According to the Shortle and Nathans method, the plasmid bearing the subtilisin gene would be nicked by a restriction enzyme that cleaves within the gene. This nick would be widened into a gap using the exonuclease action of DNA polymerase I. The resulting single-stranded gap could then be mutagenized using any one of the above mentioned mutagenizing agents.

35 Alternatively, the subtilisin gene from a Bacillus species including the natural promoter and other control sequences could be cloned into a plasmid vector containing replicons for both E. coli and B. subtilis, a selectable phenotypic marker and the M13 origin of replication for



production of single-stranded plasmid DNA upon superinfection with helper phage IR1. Single-stranded plasmid DNA containing the cloned subtilisin gene is isolated and annealed with a DNA fragment containing vector sequences but not the coding region of subtilisin, resulting in a gapped duplex molecule. Mutations are introduced into the subtilisin gene either with sodium bisulfite, nitrous acid or formic acid or by replication in a mutator strain of E. coli as described above. Since sodium bisulfite reacts exclusively with cytosine in a single-stranded DNA, the mutations created with this mutagen are restricted only to the coding regions. Reaction time and bisulfite concentration are varied in different experiments such that from one to five mutations are created per subtilisin gene on average. Incubation of 10  $\mu$ g of gapped duplex DNA in 4 M Na-bisulfite, pH. 6.0, for 9 minutes at 37°C in a reaction volume of 400  $\mu$ l, deaminates about 1% of cytosines in the single-stranded region. The coding region of mature subtilisin contains about 200 cytosines, depending on the DNA strand. Advantageously, the reaction time is varied from about 4 minutes (to produce a mutation frequency of about one in 200) to about 20 minutes (about 5 in 200).

After mutagenesis the gapped molecules are treated in vitro with DNA polymerase I (Klenow fragment) to make fully double-stranded molecules and fix the mutations. Competent E. coli are then transformed with the mutagenized DNA to produce an amplified library of mutant subtilisins. Amplified mutant libraries can also be made by growing the plasmid DNA in a Mut D strain of E. coli which increases the range for mutations due to its error prone DNA polymerase.

The mutagens nitrous acid and formic acid may also be used to produce mutant libraries. Because these chemicals are not as specific for single-stranded DNA as sodium bisulfite, the mutagenesis reactions are performed according to the following procedure. The coding portion of the subtilisin gene is cloned in M13 phage by standard methods and single stranded phage DNA prepared. The single-stranded DNA is then reacted with 1 M nitrous acid pH. 4.3 for 15-60 minutes at 23°C or 2.4 M formic acid for 1-5 minutes at 23°C. These ranges of reaction times produce a mutation frequency of from 1 in 1000 to 5 in

1000. After mutagenesis, a universal primer is annealed to the M13 DNA and duplex DNA is synthesized using the mutagenized single-stranded DNA as a template so that the coding portion of the subtilisin gene becomes fully double-stranded. At this point the coding region can be cut out of the M13 vector with restriction enzymes and ligated into an unmutagenized expression vector so that mutations occur only in the restriction fragment. (Myers et al., Science 229:242-257 (1985)).

#### 10 GENERATION OF SITE DIRECTED MUTATIONS IN THE SUBTILISIN GENE

Once the subtilisin gene has been cloned, and desirable sites for mutation identified, these mutations can be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a preferred method, a single stranded gap of DNA, bridging the subtilisin gene, is created in a vector bearing the subtilisin gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in by DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al., (1984, Biotechnology 2:646-639). According to Morinaga et al., a fragment within the gene is removed using restriction endonuclease. The vector/gene, now containing a gap, is then denatured and hybridized to a vector/gene which, instead of containing a gap, has been cleaved with another restriction endonuclease at a site outside the area involved in the gap. A single-stranded region of the gene is then available for hybridization with mutated oligonucleotides, the remaining gap is filled in by the Klenow fragment of DNA polymerase I, the insertions are ligated with T4 DNA ligase, and, after one cycle of replication, a double-stranded plasmid bearing the desired mutation is produced. The Morinaga method obviates the additional manipulation of constructing new restriction sites, and therefore facilitates the generation of mutations at multiple sites. U.S. Patent number 4,760,025, by Estell et al., issued July 26, 1988, is able to introduce oligonucleotides bearing multiple

mutations by performing minor alterations of the cassette, however, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of oligonucleotides, of various lengths, can be introduced.

5

#### EXPRESSION OF SUBTILISIN MUTANTS

According to the invention, a mutated subtilisin gene produced by methods described above, or any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector. An expression vector generally falls under the definition of a cloning vector, since an expression vector usually includes the components of a typical cloning vector, namely, an element that permits autonomous replication of the vector in a microorganism independent of the genome of the microorganism, and one or more phenotypic markers for selection purposes. An expression vector includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes. To permit the secretion of the expressed protein, nucleotides encoding a "signal sequence" may be inserted prior to the coding sequence of the gene. For expression under the direction of control sequences, a target gene to be treated according to the invention is operably linked to the control sequences in the proper reading frame. Promoter sequences that can be incorporated into plasmid vectors, and which can support the transcription of the mutant subtilisin gene, include but are not limited to the prokaryotic  $\beta$ -lactamase promoter (Villa-Kamaroff, et al., 1978, Proc. Natl. Acad. Sci. U.S.A. 75:3727-3731) and the tac promoter (DeBoer, et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:21-25). Further references can also be found in "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94.

According to one embodiment B. subtilis is transformed by an expression vector carrying the mutated DNA. If expression is to take place in a secreting microorganism such as B. subtilis a signal sequence may follow the translation initiation signal and precede the DNA sequence of interest. The signal sequence acts to transport the expression product to the cell wall where it is cleaved from the product upon secretion.

The term "control sequences" as defined above is intended to include a signal sequence, when it is present.

#### EXAMPLES

### 5 SITE SPECIFIC MUTATION OF THE SUBTILISIN GENE GENERATES MUTANTS WITH USEFUL CHEMICAL CHARACTERISTICS

#### MATERIALS AND METHODS

### 10 BACTERIAL STRAINS

B. subtilis 309 and 147 are variants of Bacillus lentus, deposited with the NCIB and accorded the accession numbers NCIB 10147 and NCIB 10309, and described in U.S. Patent No. 3,723,250, issued March 27, 1973, and incorporated in its  
15 entirety by reference herein.

B. subtilis DN 497 is described in U.S. Serial No. 039,298 filed April 17, 1987, corresponding to EP Publ. No. 242 220, also incorporated by reference herein, and is an aro<sup>+</sup> transformant of RUB 200 with chromosomal DNA from SL 438, a  
20 sporulation and protease deficient strain obtained from Dr. Kim Hardy of Biogen.

E. coli MC 1000 r<sup>m+</sup> (Casadaban, M.J. and Cohen, S.N. (1980), J. Mol. Biol. 138: 179-207), was made r<sup>m+</sup> by conventional methods and is also described in U.S. Serial No.  
25 039,298.

B. subtilis DB105 is described in: Kawamura, F., Doi, R.H. (1984), Construction of a Bacillus subtilis double mutant deficient in extracellular alkaline and neutral proteases, J. Bacteriol. 160 (2), 442- 444.

30

#### PLASMIDS

pSX50 (described in U.S. patent application serial No. 039,298, and incorporated by reference herein) is a derivative of plasmid pDN 1050 comprising the promoter-operator P<sub>1</sub>O<sub>1</sub>, the  
35 B. pumilus xyn B gene and the B. subtilis xyl R gene.

pSX62 (described in U.S. patent application serial No. 039,298, supra) is a derivative of pSX52 (ibid), which comprises a fusion gene between the calf prochymosin gene and the B. pumilus xyn B gene inserted into pSX50 (supra). pSX62

was generated by inserting the E. coli rrn B terminator into pSX52 behind the prochymosin gene.

pSX65 (described in U.S. patent application serial No. 039,298, supra) is a derivative of plasmid pDN 1050, comprising the promotor-operator P<sub>2</sub>O<sub>2</sub>, the B. pumilus xyn B gene, and the B. subtilis xyl R gene.

pSX88 (described in unpublished International Patent Application no. PCT/DK 88/00002 (NOVO INDUSTRI A/S)) is a derivative of pSX50 comprising the subtilisin 309 gene.

pSX92 was produced by cloning the subtilisin 309 into plasmid pSX62 (supra) cut at Cla I and Hind III, and Cla I filled prior to the insertion of the fragments DraI-NheI and NheI-Hind III from the cloned subtilisin 309 gene.

pSX93, shown in Figure 3, is pUC13 (Vieira and Messing, 1982, Gene 19:259-268) comprising a 0.7kb XbaI-Hind III fragment of the subtilisin 309 gene including the terminator inserted in a polylinker sequence.

pSX119 (described in unpublished International Patent Application no. PCT/DK 88/00002 (NOVO INDUSTRI A/S)) is pUC13 harbouring an EcoRI-XbaI fragment of the subtilisin 309 gene inserted into the polylinker.

pSX120 is a plasmid where the HpaI-HindIII fragment with the subtilisin 309 gene from pSX88 is inserted into EcoRV-HindIII on pDN 1681, in a way whereby the protease gene is expressed by the amy M and amy Q promoters. pDN 1681 is obtained from pDN 1380 (Diderichsen, B. and Christiansen, L.: 1988, FEMS Microbiology Letters 56: 53-60) with an inserted 2.85 bp ClaI fragment from B. amyloliquefaciens carrying the amy Q gene with promotor (Takkinen et al.: 1983, J. Biol. Chem. 258: 1007ff.)

pUC13 is described in: Vieira, J. and Messing, J.: 1982, Gene 19: 259-268.

pUC19 is described in: Yanisch-Perron, C. and Vieira, J. Messing, J., 1985, Gene 33:103-119.

pUB110 is described in: Lacey, R.W., Chopra, J. (1974), Genetic studies of a multiresistant strain of Staphylococcus aureus, J. Med. Microbiol. 7, 285-297, and in: Zyprian, E., Matzura, H. (1986), Characterization of signals promoting gene expression on the Staphylococcus aureus plasmid pUB110 and

development of a Gram-positive expression vector system, DNA  
5 (3), 219-225.

### GENES

5           The genes for the various subtilisins were obtained as  
referenced in the literature mentioned above. In particular  
the genes for the subtilisin 309 and 147 enzymes were obtained  
as described in unpublished international patent application  
no. PCT/DK 88/00002 (NOVO INDUSTRI A/S), which is hereby  
10 incorporated by reference in its entirety.

### SUBTILISIN CARLSBERG GENE CONSTRUCTION

A synthetic gene was designed based on the coding  
sequence of the mature subtilisin Carlsberg protease and its  
15 transcription terminator (Jacobs, M. Eliasson, M., Uhlen, M.,  
Flock, J.-I. (1985), Cloning, sequencing and expression of  
subtilisin Carlsberg from Bacillus licheniformis. Nucleic  
Acids Res. 13 (24), 8913-8926), linked to the pre and pro  
coding sequences of the subtilisin BPN' protease (Wells, J.A.,  
20 Ferrari, E., Henner, D.J., Estell, D.A., Chen, E.Y. (1983),  
Cloning, sequencing and secretion of Bacillus amyloliquefa-  
ciens subtilisin in Bacillus subtilis, Nuclei Acids Res. 11  
(22), 7911-7925). The gene was subdivided into seven fragments  
in length ranging from 127 to 313 basepairs, each fragment  
25 built up from chemically synthesized oligos of 16 to 77  
nucleotides. The overlap between the oligos of the two strands  
was optimised in order to facilitate a one step annealing of  
each fragment (Mullenbach, G.T., Tabrizi, A., Blacher, R.W.,  
Steimer, K.S. (1986), Chemical synthesis and expression in Yeast  
30 of a gene encoding connective tissue activating peptide-III,  
J. Biol. Chem. 261 (2), 719-722). Each fragment was assembled  
and cloned in an E. Coli cloning and sequencing vector.  
Sequence analysis of these cloned fragments was performed to  
confirm the correctness of the sequence of each fragment. Then  
35 all of the fragments were assembled and cloned in the vector  
pUB110 (Lacey, R.W., Chopra, J. (1974), Genetic studies of a  
multiresistant strain of Staphylococcus aureus, J.  
Med. Microbiol. 7, 285-297) and brought into B. subtilis DB105  
(Kawamura, F., Doi, R.H. (1984), Construction of a Bacillus

subtilis double mutant deficient in extracellular alkaline and neutral proteases, J.Bacteriol. 160 (2), 442-444). Transcription of the gene was initiated by the HpaII promotor of the pUB110 plasmid vector (Zyprian, E., Matzura, H. (1986), Characterization of signals promoting gene expression on the Staphylococcus aureus plasmid pUB110 and development of a Gram-positive expression vector system, DNA 5 (3), 219-225). In the process of the gene construction it turned out that the longest fragment (#5; 313 basepairs long) needed further fragmentation (fragments #8 and #9) in order to avoid problems with the assembly of this rather long fragment.

The amino acid sequence deduced from the nucleotide sequence differs from the earlier published subtilisin Carlsberg sequence at positions 129, 157, 161 and 212 (Smith, E.L., DeLange, R.J., Evans, W.H., Landon, W., Markland, F.S. (1968), Subtilisin Carlsberg V. The complete sequence: comparison with subtilisin BPN'; evolutionary relationships., J.Biol.Chem. 243 (9), 2184-2191). A fifth alteration reported by Jacobs et al (1985) could not be confirmed in the clone of the Carlsberg gene described here.

#### COMPUTATION OF ISOELECTRIC POINT ( $pI_0$ )

The calculation of the isoelectric point of the subtilisin 309 wild type enzyme (S000) is exemplified below in order to demonstrate the procedure used. The same procedure is of course applicable to the computation of any enzyme, whether it being a mutant enzyme or not.

pK values were assigned to each potentially charged amino acid residue (Tyr, Asp, Glu, Cys, Arg, His, Lys, N-terminal, C-terminal,  $Ca^{2+}$ ). In this case the environment was taken into consideration, whereby different pK values are used for the same amino acid residue dependent on its neighbours. The assigned values are indicated in Table II.

Then the ratio of the occurrence of an amino acid residue at a given pH in charged or uncharged form (charged/uncharged, C/U(i)) was calculated for both negative and positive charge, by using the formulae Ia and Ib, respectively. In Table II this ratio is only indicated for pH equal to  $pI_0$ .

Subsequently the relative charge,  $Q_r(i)$ , or charge contribution allocated to each charged residue was calculated by using the formulae IIa and IIb:

The pH value where the sum of all the charge contributions from the charged residues is equal to zero was found by iteration.

Table II

Calculation of isoelectric point for: S000 Subtilisin 309

10

Residue	Number of C/U(i)*		$Q_r(i)$	$Q_r(i)$		$Q_r(i)$
	pK	Residue		pH 8.3	pH 10.0	pH=pI <sub>0</sub>
Tyr	9.9	3	2.51E-02	-0.07	-1.67	-1.77
15 Tyr	11.6	2	5.01E-04	0.00	-0.05	-0.06
Tyr	12.5	2	6.31E-05	0.00	-0.01	-0.01
Asp	3.5	5	6.31E+04	-5.00	-5.00	-5.00
Glu	4	5	2.00E+04	-5.00	-5.00	-5.00
C-term(Arg)	3	1	2.00E+05	-1.00	-1.00	-1.00
20 Cys	9.3	0	1.00E-01	0.00	0.00	0.00
Arg	12.8	8	3.16E+04	8.00	7.99	7.99
His	6.4	7	1.26E-02	0.09	0.00	0.00
Lys	10	5	5.01E+01	4.90	2.50	2.34
Calcium	20	1.25	5.01E+11	2.50	2.50	2.50
25 N-term(Ala)	8	1	5.01E-01	0.33	0.01	0.01
Net charge				4.75	0.27	0.0
The calculated isoelectric point is				10.06		

30 \* E-02 = 10<sup>-2</sup>

As indicated above and in Table II the pK value assigned to each amino acid was different taking local variations in the environment into consideration. This only results in an enhanced precision in the calculation, but experience has shown that constant estimated pK values are helpful in showing in what direction the pI<sub>0</sub> for a given mutant enzyme will move in comparison to the pI<sub>0</sub> of the parent enzyme. This is indi-



cated in Table III, where  $pI_o$  values for estimated  $pK$  values are indicated.

In order to compare various enzymes and mutant enzymes washing tests described in detail below have been performed.

5 In Table III below results from these tests using parent enzyme and mutant enzymes from subtilisin 309 (designated S000, etc.) and subtilisin Carlsberg (designated C000, etc.) have been tabulated in order to demonstrate the correlation between  $pI_o$  and wash performance at different pH values of the wash liquor  
10 used. In the washing tests a low salt liquid detergent formulation of pH 8.3 according to detergent example D7, and a normal salt powder detergent of pH 10.2 according to detergent example D2 were used.

In Table III the results are indicated as relative  
15 results compared to the wild type enzymes (S000 and C000, respectively). Also, calculated and observed  $pI_o$ s for the enzymes are indicated.

Table III

Comparative washing tests at different pH values

	Mutant	$pI_0$		<u>Improvement Factor</u>	
		calculated	observed	Detergent pH	
				8.3	10.2
5	S000	10.02	9.7	1	1
	S001	9.86	9.4	2.2	1
	S003	9.86	9.4	2.0	1
	S004	9.68	9.1	3.9	1
10	S005	9.71	9.1	1.5	1
	S012	9.09	8.8	5.0	0.6
	S019	9.09	8.5	5.8	0.6
	S020	6.71	7.9	8.8	0.5
	S021	9.85	-	1.8	0.7
15	S022	8.07	-	9.0	0.3
	S023	8.05	-	9.8	0.2
	S024	6.86	-	9.0	0.2
	S025	8.94	-	6.9	0.6
	S027	10.28	-	0.4	1.0
20	S028	10.28	-	0.9	1.0
	S031	10.53	-	0.4	0.7
	S032	10.28	-	0.7	-
	S033	10.28	-	0.4	-
	S035	8.07	-	8.0	0.6
25	S201	9.85	-	2.0	0.7
	S202	9.62	-	4.3	0.9
	S203	9.27	-	9.0	0.5
	C000	8.87	-	1	1
	C001	9.38	-	0.2	1.5
30	C002	9.38	-	0.8	1.9
	C003	9.38	-	0.4	1.1
	C004	9.64	-	0.2	1.8
	C008	9.38	-	0.2	1.5

35 From Table III it is seen that shifting the  $pI_0$  to lower values (S-series) provides for an improvement in wash performance at low pH (pH=8.3), whereas an upward shift in  $pI_0$  (C-series) provides for an improvement in wash performance at high pH (pH=10.2).

The concept of isoelectric point has thus been found to be very useful in selecting the positions of the amino acids in the parent enzyme should be changed.

- 5 It has generally been found that mutations should be performed in codons corresponding to amino acids situated at or near to the surface of the enzyme molecule thereby retaining the internal structure of the parent enzyme as much as possible.

10

#### PURIFICATION OF SUBTILISINS

The procedure relates to a typical purification of a 10 litre scale fermentation of the Subtilisin 147 enzyme, the Subtilisin 309 enzyme or mutants thereof.

- 15 Approximately 8 litres of fermentation broth were centrifuged at 5000 rpm for 35 minutes in 1 litre beakers. The supernatants were adjusted to pH 6.5 using 10% acetic acid and filtered on Seitz Supra S100 filter plates.

The filtrates were concentrated to approximately 400 ml using an Amicon CH2A UF unit equipped with an Amicon S1Y10 UF cartridge. The UF concentrate was centrifuged and filtered prior to absorption at room temperature on a Bacitracin affinity column at pH 7. The protease was eluted from the Bacitracin column at room temperature using 25% 2-propanol and 25 1 M sodium chloride in a buffer solution with 0.01 dimethylglutaric acid, 0.1 M boric acid and 0.002 M calcium chloride adjusted to pH 7.

The fractions with protease activity from the Bacitracin purification step were combined and applied to a 750 ml Sephadex G25 column (5 cm dia.) equilibrated with a buffer containing 0.01 dimethylglutaric acid, 0.2 M boric acid and 0.002 m calcium chloride adjusted to pH 6.5.

Fractions with proteolytic activity from the Sephadex G25 column were combined and applied to a 150 ml CM Sepharose CL 6B cation exchange column (5 cm dia.) equilibrated with a buffer containing 0.01 M dimethylglutaric acid, 0.2 M boric acid, and 0.002 M calcium chloride adjusted to pH 6.5.

The protease was eluted using a linear gradient of 0-0.1 M sodium chloride in 2 litres of the same buffer (0-0.2 M sodium chloride in case of sub 147).

In a final purification step protease containing 5 fractions from the CM Sepharose column were combined and concentrated in an Amicon ultrafiltration cell equipped with a GR81PP membrane (from the Danish Sugar Factories Inc.).

Subtilisin 309 and mutants

Gly 195 Glu (G195E (S001)):  
 10 Arg 170 Tyr (R170Y (S003)):  
 Arg 170 Tyr + Gly 195 Glu (R170Y+G195E (S004)):  
 Lys 251 Glu (K251E (S005)):  
 His 120 Asp (H120D (S006)):  
 Arg 170 Tyr + Gly 195 Glu + Lys 251 Glu  
 15 (R170Y+G195E+K251E (S012)):  
 Lys 235 Leu (K235L (S015)):  
 His 120 Asp + Gly 195 Glu + Lys 235 Leu (H120D+G195E+  
 K235L (S017)):  
 His 120 Asp + Arg 170 Tyr + Gly 195 Glu + Lys 235  
 20 Leu (H120D+R170Y+G195E+K235L (S019)) :  
 His 120 Asp + Arg 170 Tyr + Gly 195 Glu + Lys 235 Leu  
 + Lys 251 Glu (H120D+R170Y+G195E+K235L+K251E (S020):

were purified by this procedure.

25

#### PURIFICATION OF (MUTANT) SUBTILISIN CARLSBERG PROTEASES

Fermentation media were either directly applied on a bacitracin affinity column (5 cm diam \* 15 cm; equilibrated with 10 mM Tris/HCl buffer pH 7.9; flow rate approx. 500 ml/h)  
 30 or concentrated to 500 ml by means of a Nephross Andante H.F. dialyzer (Organon Technika) using a back pressure of 10-12 p.s.i. and demineralized water in the outer circuit. In the latter case the protease was precipitated from the concentrate by adding 600 g/l ammonium sulphate. The precipitate was  
 35 collected by means of centrifugation and redissolved in approx. 500 ml demineralized water. The ammonium sulphate was removed from the protease solution using the same dialyzer as described above. The final volume was approx. 300 ml, while the pH was adjusted to pH 6.0. The protease was eluted from the bacitracin

columns (mentioned above) using a 10 mM Tris buffer (pH 7.9) containing 2.7 M NaCl and 18% isopropanol.

After dialysis of bacitracin-purified or concentrated protease material further purification was accomplished by application on a CM-Trisacryl ion exchange column (5 cm. diam \* 15 cm; equilibrated with 0.03M sodium phosphate pH 6.0) using a flow rate of 200 ml/h. The protease was eluted from the column with a linear gradient from 0 to 0.3 M NaCl (2 \* 500 ml) in the phosphate buffer. Fractions containing protease activity were pooled and stored at -20°C in the presence of buffer salts after freeze-drying.

#### OLIGONUCLEOTIDE SYNTHESIS

All mismatch primers were synthesized on an Applied Biosystems 380 A DNA synthesizer and purified by polyacrylamide gel electrophoresis (PAGE).

#### ASSAY FOR PROTEOLYTIC ACTIVITY

The proteolytic activity of the mutant enzymes was assayed in order to determine how far the catalytic activity of the enzyme was retained. The determinations were performed by the dimethyl casein (DMC) method described in NOVO Publication AF 220-gb (or later editions), available from Novo-Nordisk a/s, Bagsværd, Denmark, which publication is hereby included by reference.

#### ASSAYS FOR WASH PERFORMANCE

A:

Test cloths (2.2cm x 2.2cm), approximately 0.1 g) were produced by passing desized cotton (100% cotton, DS 71) cloth through the vessel in a Mathis Washing and Drying Unit type TH (Werner Mathis AG, Zurich, Switzerland) containing grass juice.

Finally the cloth was dried in a strong air stream at room temperature, stored at room temperature for 3 weeks, and subsequently kept at -18° C prior to use.

All tests were performed in a model miniwash system. In this system 6 test cloths are washed in a 150 ml beaker containing 60 ml of detergent solution. The beakers are kept in a thermostat water bath at 30° C with magnetic stirring.

As detergent the following standard liquid detergent was used:

	AE, Berol 160	15%
5	LAS, Nasa 1169/P	10%
	Coconut fatty acid	9%
	Oleic acid	1%
	Triethanolamine	9%
	Glycerol	10.5%
10	Ethanol	1.5%
	Tri·Na·Citrat·2H <sub>2</sub> O	8%
	CaCl·2H <sub>2</sub> O	0.1%
	NaOH	1%
	Water from LAS	23.3%
15	Water from glycerol	1.5%
	Water added	34.9%

The percentages given are the percentage of active content.

pH was adjusted with 1 N NaOH to 8.14. The water used was ca. 6° dH (German Hardness).

Tests were performed at enzyme concentrations of: 0, 1.0 mg enzyme protein/l and 10.0 mg enzyme protein/l, and two independent sets of tests were performed for each of the mutants. The results shown in the following are means of these tests.

The washings were performed for 60 minutes, and subsequent to the washing the cloths were flushed in running tap-water for 25 minutes in a bucket.

The cloths were then air-dried overnight (protected against daylight) and the remission, R, determined on an ELREPHO 2000 photometer from Datacolor S.A., Dietlikon, Switzerland at 460 nm.

As a measure of the wash performance differential remission, delta R, was used being equal to the remission after wash with enzyme added minus the remission after wash with no enzyme added.

B:

The wash performance of various mutants was tested against grass juice stained cotton cloths according to the method described above.

2.0 g/l of a commercial US liquid detergent was used.

5 The detergent was dissolved in a 0,005 M ethanolamine buffer in ion-exchanged water. pH was adjusted to pH 8.0, 9.0, 10.0 and 11.0 respectively with NaOH/HCl.

The temperature was kept at 30° C isothermic for 10 min.

10 The mutants were dosed at 0.25 mg enzyme protein/l each.

C:

Washing tests using the detergent compositions  
15 exemplified in the detergent examples below were performed in a mini washer utilizing cotton based test cloths containing pigments, fat, and protein (casein). The conditions were:

a) 2 g/l detergent D3 in 6°fH (French hardness) water at pH 8.3, or

20 b) 5 g/l detergent D2 in 15°fH water at pH 10.2.

After rinsing and drying reflection at 460 nm was measured.

The improvement factor was calculated from a dose-response curve, and relates to the amount of enzyme needed for  
25 obtaining a given delta R value in comparison to the wild type enzyme in question (S000 and C000), meaning that an improvement factor of 2 indicates that only half the amount of enzyme is needed to obtain the same delta R value.

The results of these tests are shown in Table III  
30 above.

D:

Experimental tests of lipase stability were carried out for example using the following materials:

35

1 LU/ml Pseudomonas cepacia lipase was incubated in wash liquor of each of two types, O and W (described below). Aliquots were taken at intervals and tested for lipase activity. Parallel incubations were carried out without

protease or with protease of various types as noted below, to test the effect of the protease on the retention of lipase activity. Wild-type proteases were tested at 20 GU/ml, mutated proteases were tested at 0.5 microgram/ml.

5

#### DETERGENT COMPOSITIONS COMPRISING ENZYME VARIANTS

##### 10 Detergent D1:

A detergent powder according to an embodiment of the invention containing phosphate builder is formulated to contain: total active detergent about 16%, anionic detergent about 9%, nonionic detergent about 6%, phosphate-containing  
15 builder about 20%, acrylic or equivalent polymer about 3.5%, (alternatively down to about 2%), perborate bleach precursor about 6-18%, alternatively about 15-20%, amino-containing bleach activator about 2%, silicate or other structurant about 3.5%, alternatively up to about 8%, enzyme of about 8 glycine  
20 units/mg activity, with alkali to adjust to desired pH in use, and neutral inorganic salt, and enzymes (about 0.5% each enzyme).

The anionic detergent is a mixture of sodium dodecyl-benzene sulphonate, alternatively sodium linear alkyl-ben-  
25 zene-sulphonate, 6%, and primary alkyl sulphate 3%. The nonionic detergent is an ethoxylate of an approx. C13-C15 primary alcohol with 7 ethoxylate residues per mole. The phosphate builder is sodium tripolyphosphate. The polymer is polyacrylic acid, alternatively acrylic/maleic copolymer. The  
30 perborate bleach precursor is sodium tetraborate tetrahydrate or monohydrate. The activator is tetra-acetyl-ethylene-diamine. The structurant is sodium silicate. The neutral inorganic salt is sodium sulphate.

The enzymes comprise protease according to Mutant  
35 S001, alternatively protease S003, S004, S005, C001, C002, C003, C004, C005, C008, S015, S017, S021, S226, S223, S224, or S225.

Detergent D1a:



A detergent powder according to an embodiment of the invention containing phosphate builder is formulated to contain: total active detergent about 15%, anionic detergent about 7%, nonionic detergent about 6%, phosphate-containing builder about 25%, acrylic or equivalent polymer about 0.5%, perborate bleach precursor about 10%, amino-containing bleach activator about 2%, silicate or other structurant about 6%, protease enzyme of about 8 glycine units/mg grade, with alkali to adjust to desired pH in use, and neutral inorganic salt, and enzymes (about 0.5% each enzyme).

The anionic detergent is sodium linear alkyl-benzene-sulphonate. The nonionic detergent is an ethoxylate of an approx. C13-C15 primary alcohol with 7 ethoxylate residues per mole or a mixture of this with the corresponding alcohol ethoxylated to the extent of 3 residues per mole. The phosphate builder is sodium tripolyphosphate. The perborate or peracid bleach precursor is sodium tetraborate tetrahydrate. The activator is tetra-acetyl-ethylene-diamine. The structurant is sodium silicate. The neutral inorganic salt is sodium sulphate. The enzymes comprise protease according to Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S015, S017, S021, S226.

#### Detergent D2:

A detergent powder according to an embodiment of the invention containing zeolite builder is formulated to contain: total active detergent about 16%, anionic detergent about 9%, nonionic detergent about 6%, zeolite-containing builder about 20%, acrylic or equivalent polymer about 3.5%, perborate bleach precursor about 6-18%, amino-containing bleach activator about 2%, silicate or other structurant about 3.5%, alternatively down to about 2.5%, enzyme of about 8 (alternatively about 15) glycine units/mg grade, with alkali to adjust to desired pH in use, and neutral inorganic salt, and enzymes (about 0.5% each enzyme).

The anionic detergent is a mixture of sodium dodecyl-benzene sulphonate, alternatively sodium linear alkyl-benzene-sulphonate, 6% and primary alkyl sulphate 3%. The nonionic detergent is an ethoxylate of an approx. C13-C15 primary

alcohol with 7 ethoxylate residues per mole. The zeolite builder is type A zeolite. The polymer is polyacrylic acid. The perborate bleach precursor is sodium tetraborate tetrahydrate or monohydrate. The activator is tetraacetyl-ethylenediamine.  
5 The structurant is sodium silicate. The neutral inorganic salt is sodium sulphate. The enzymes comprise protease according to Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S015, S017, S021, S226.

#### 10 Detergent D2a:

A detergent powder according to an embodiment of the invention containing zeolite builder is formulated to contain: total active detergent about 14%, anionic detergent about 7%, nonionic detergent about 7%, zeolite-containing builder about  
15 25%, acrylic or equivalent polymer about 3%, perborate or peracid bleach precursor about 10%, amino-containing bleach activator about 2%, silicate or other structurant about 0.5%, enzyme of about 6 glycine units/mg grade, with alkali to adjust to desired pH in use, and neutral inorganic salt, and enzymes  
20 (about 0.5% each enzyme).

The anionic detergent is sodium linear alkyl-benzene-sulphonate, the nonionic detergent is a mixture of ethoxylates of an approx. C13-C15 primary alcohol with 7 and 3 ethoxylate residues respectively per mole. The zeolite builder  
25 is type A zeolite. The polymer is an acrylic/maleic copolymer. The perborate bleach precursor is sodium tetraborate monohydrate. The activator is tetra-acetyl-ethylene-diamine. The structurant is sodium silicate. The neutral inorganic salt is sodium sulphate. The enzymes comprise protease according  
30 to Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S015, S017, S021, S226.

#### Detergent D3:

An aqueous detergent liquid according to an embodiment of the invention is formulated to contain: Dodecylbenzene-  
35 -sulphonic acid 16%, C12-C15 linear alcohol condensed with 7 mol/mol ethylene oxide 7%, monoethanolamine 2%, citric acid 6.5%, sodium xylenesulphonate 6%, sodium hydroxide about 4.1%, protease 0.5%, minors and water to 100%. The pH is adjusted to

a value between 9 and 10. The enzyme is a protease according to Mutant S020, alternatively S019, S012, S004, S001, S003, S005, S015, S017, S021, S022, S025, S035, S201, S223, S224, S225, S226 or S235.

5

Detergent D4:

A nonaqueous detergent liquid according to an embodiment of the invention is formulated using 38.5% C13-C15 linear primary alcohol alkoxylated with 4.9 mol/mol ethylene  
10 oxide and 2.7 mol/mol propylene oxide, 5% triacetin, 30% sodium triphosphate, 4% soda ash, 15.5% sodium perborate monohydrate containing a minor proportion of oxoborate, 4% TAED, 0.25% EDTA of which 0.1% as phosphonic acid, Aerosil 0.6%, SCMC 1%, and 0.6% protease. The pH is adjusted to a value between 9 and 10,  
15 e.g. about 9.8. The enzyme comprises protease according to Mutant S001, alternatively S003 or S004, S021, S035, S201, S225, S226, or S235.

Detergent D5:

20 A detergent powder according to an embodiment of the invention is formulated in the form of a granulate having a bulk density of at least 600 g/l, containing about 20% by weight surfactant of which about 10% is sodium dodecylbenzene sulphonate, and the remainder is a mixture of Synperonic A7 and  
25 Synperonic A3 (about 5.5% to 4.5%), and zero neutral inorganic salt (e.g. sodium sulphate), plus phosphate builder about 33%, sodium perborate tetrahydrate about 16%, TAED activator about 4.5%, sodium silicate about 6%, and minors including sodium carbonate about 2%, and moisture content about 10%. Enzymes  
30 (about 0.5% each enzyme) are included. The enzyme comprises protease according to Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S223, S224, S225, S226 or S235.

35 Detergent D6:

A detergent powder according to an embodiment of the invention is formulated in the form of a granulate having a bulk density of at least 600 g/l, alternatively about 550 g/l, containing about 20%, alternatively down to about 16%, by

weight surfactant of which about 9%, alternatively about 7%, is sodium dodecylbenzene sulphonate, alternatively sodium linear alkyl benzene sulphonate, and the remainder is a mixture of Synperonic A7 and Synperonic A3 (or similar ethoxylates) (respectively about 5% & 6%, alternatively about 4% and 7%), and zero neutral inorganic salt (e.g. sodium sulphate), plus zeolite builder about 30%, alternatively about 25%, sodium perborate tetrahydrate, alternatively monohydrate, about 14% or 15%, TAED activator about 3.6%, and minors including sodium carbonate about 9%, or up to 15%, Dequest® 2047 about 0.7%, and moisture content about 10%. Enzymes (about 0.5% each enzyme, or about 0.2% lipase and about 0.7% protease) are included. The enzyme comprises protease according to Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S223, S224, S225, S226, or S235.

#### Detergent D6a:

A detergent powder according to an embodiment of the invention is formulated in the form of a granulate having a bulk density of at least 600 g/l, containing about 15% by weight surfactant of which about 7% is sodium linear alkyl benzene sulphonate, 2% primary alcohol sulphate, and the remainder Synperonic A7 or similar ethoxylate, and zero neutral inorganic salt (e.g. sodium sulphate), plus zeolite builder about 22%, sodium perborate tetrahydrate about 15%, TAED activator about 7%, and minors including sodium carbonate about 15%, Dequest® 2047 about 0.7%, and moisture content about 10%. Enzymes (about 1.2%) include protease Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S223, S224, S225, S226 or S235.

#### Detergent D7:

A detergent powder according to an embodiment of the invention is formulated to contain: Dodecylbenzenesulphonic acid 6%, C12-C15 linear alcohol condensed with 7 mol/mol ethylene oxide 5%, fatty acid soap 3%, Sokolan® CP5 polymer 3%, zeolite A 22%, sodium carbonate 10%, sodium sulphate 17%, clay particles 8%, sodium perborate tetrahydrate 13%, tetraacetyl-ethylenediamine 2%, protease 0.5%, minors and water to

100%. The pH is adjusted to a value between 9 and 10. The enzyme comprises protease Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S223, S224, S225, S226 or S235.

5

#### Detergent D8:

A detergent (soap) bar according to an embodiment of the invention is formulated as follows: soap based on pansaponified 82% tallow, 18% coconut oil, neutralised with 0.15% orthophosphoric acid, mixed with protease (about 8 GU/mg of the bar composition) and mixed with sodium formate 2%, borax 2%, propylene glycol 2% and sodium sulphate 1%, is then plodded on a soap production line. The enzyme comprises Mutant S001, alternatively S003, S004, S005, C001, C002, C003, C004, C005, C008, S021, S025, S035, S201, S202, S223, S224, S225, S226 or S235.

#### Detergent D9:

Structured liquid detergents can for example contain, in addition to a protease as described herein, 2-15% nonionic surfactant, 5-40% total surfactant, comprising nonionic and optionally anionic surfactant, 5-35% phosphate-containing or non-phosphate containing builder, 0.2-0.8% polymeric thickener, e.g. cross-linked acrylic polymer with m.w. over 106, at least 10% sodium silicate, e.g. as neutral waterglass, alkali (e.g. potassium-containing alkali) to adjust to desired pH, preferably in the range 9-10 or upwards, e.g. above pH 11, with a ratio sodium cation: silicate anion (as free silica) (by weight) less than 0.7:1, and viscosity of 0.3-30 Pas (at 20°C and 20s-1).

Suitable examples contain about 5% nonionic surfactant C13-15 alcohol alkoxylated with about 5 EO groups per mole and with about 2.7 PO groups per mole, 15-23% neutral waterglass with 3.5 weight ratio between silica and sodium oxide, 13-19% KOH, 8-23% STPP, 0-11% sodium carbonate, 0.5% Carbopol® 941.

Protease (e.g. 0.5%) includes Mutant S001, alternatively S021, S025, S035, S201, S202, S223, S224, S225, S226 or S235.

## Detergent D10

A structured, viscous, aqueous liquid detergent suitable for laundry use is formulated as follows (% by weight):

	Citric acid	2.5
	Borax (10aq)	4
	NaOH	2
	Glycerol	5
10	C14-C15 Linear alkyl-benzene-sulphonate, or C14-15 primary alcohol sulphate	6.5
	Synperonic A3	
	Nonionic C12-C15 3EO	1.2
15	Synperonic A7	
	Nonionic C12-C15 7EO	3.6
	Zeolite	20
	Protease	0.5
	Amylase (Termamyl® 300LDX)	0.2
20	minors and water	to 100%

The pH can be adjusted to a value between 9 and 10. The enzyme comprises protease Mutant S020, alternatively S019, S012, S004, S001, S003, S005, S021, S035, S201, S223, S224, S225, S226 or S235.

## Detergent D11

An isotropic aqueous liquid detergent suitable for laundry use is formulated as follows (% by weight):

	Citric acid	2
	Boric acid	1
	NaOH	3
	KOH	4.5
5	Glycerol	10
	Ethanol	6.5
	Nonionic surfactant	
	(C12-alcohol 6.5 EO	
	ethoxylate groups/mol)	
10	or sodium primary alcohol sulphate	10
	Oleic acid	16
	Coconut oil (C12) soap	11
	Protease	0.5
	minors and water	to 100%

15

The pH can be adjusted to a value between 9 and 10.  
The enzyme comprises protease Mutant S020, alternatively S019, S012, S004, S001, S003, S005, S021, S025, S035, S201, S223, S224, S225, S226 or S235.

20

#### Detergent D12

An aqueous liquid detergent composition is formulated to contain:

	sodium alkyl-benzene-sulphonate	14.5
25	C18 sodium soap	2
	Nonionic detergent (C12-15 6EO)	9
	Fatty acid (oleic acid)	4.5
	sodium alkenyl succinate	11
	propanediol	1.5
30	ethanol	3.6
	sodium citrate	3.2
	Complexing agent e.g. Dequest 2060	0.7
	Protease	0.5
	Amylase	0.1
35	Sodium chloride	0.5
	minors and water	to 100%

The pH can be adjusted to a value between 9 and 10.  
The enzyme comprises protease Mutant S020, alternatively S019,

S012, S004, S001, S003, S005, S021, S025, S035, S201, S202, S223, S224, S225, S226 or S235.

#### Detergent D13

5 An aqueous liquid detergent composition is formulated to contain:

	sodium alkyl-benzene-sulphonate	8
	nonionic detergent 6.5EO	10
	Oleic diethylamide	10
10	Fatty acid (C12/C18 75:25)	18
	sodium citrate	1
	triethanolamine	5
	propanol	7
	ethanol	5
15	Dequest 2060	0.5
	Protease	0.5
	Amylase	0.1
	minors and water	to 100%

20 The pH can be adjusted to a value between 9 and 10.  
The enzyme comprises protease Mutant S020, alternatively S019, S012, S004, S001, S003, S005, S021, S025, S035, S201, S202, S223, S224, S225, S226 or S235.

#### 25 Detergent D14

A non-aqueous liquid detergent composition is formulated to contain (% by weight):



	Liquid nonionic detergent (C10-12, 6.2EO)	41%
	triacetin	5
	linear alkylbenzenesulphonic acid	6
	magnesium oxide stabiliser	1
5	Sodium carbonate builder/base	18
	Calcium carbonate builder	8
	bleach activator TAED	3.5
	bleach precursor perborate monohydrate	10.5
	partly-hydrophobic silica	2
10	protease	0.4
	lipase (Lipolase®)	3
	minors or additional	
	liquid nonionic surfactant (no water)	to 100%

15 In formulating this composition, the liquid nonionic surfactant and triacetin are added first, followed by the magnesium oxide, then the other ingredients except enzyme. The mixture is milled in a colloid mill and cooled, and finally the enzyme(s) and any other heat-sensitive minors are added.

20 The enzyme comprises protease Mutant S020, alternatively S019, S012, S004, S001, S003, S005, S021, S025, S035, S201, S202, S223, S224, S225, S226 or S235.

Also usable are any one of the detergent formulations  
25 described and exemplified in EP 0 342 177.

Also usable are detergent formulations exemplified in EP 0 342 177, in conjunction with mutants as for detergent D3.

#### RESULTS

30

#### GENERATION OF SITE-SPECIFIC MUTATIONS OF THE SUBTILISIN 309 GENE

Site specific mutations were performed by the method  
35 of Morinaga et al. (Biotechnology, supra). The following oligonucleotides were used for introducing the mutations:

a) Gly 195 Glu (G195E (S001)):

A 27-mer mismatch primer, Nor-237, which also generates a novel SacI restriction site:

64

5' CACAGTATGGGCGCAGGGCTTGACATTGTCGCACCAGG 3'  
 Nor-237 5' GTATGGCGCAGAGCTCGACATTTGTCGC 3'

SacI

5 b) Arg 170 Tyr (R170Y (S003)):

A 25-mer mismatch primer, Nor-577, which destroys a HaeIII site:

HaeIII

5' GCTATCCGGCCCGTTATGCGAACGC 3'  
 10 Nor-577 3' CGATAGGCCCGTATAATACGCTTGCG 5'

c) His 120 Asp (H120D (S006)):

A 32-mer mismatch primer, Nor-735, which destroys a SphI site:

SphI

15 5' AGGGAACAATGGCATGCACGTTGCTAATTTGA 3'  
 Nor-735 5' AGGGAACAATGGCATGGACGTTGCTAATTTGA 3'

d) Lys 251 Glu (K251E (S005)):

20 A 32-mer mismatch primer, Nor-736, which generates a XhoI site:

5' CAAATCCGCAATCATCTAAAGAATACGGCAAC 3'  
 Nor-736 5' CAAATCCGCAATCATCTCGAGAATACGGCAAC 3'

XhoI

25

e) Lys 235 Leu (K235L (S015)):

A 24-mer mismatch primer, Nor77-856, which generates a PstI site:

5' GCCCTTGTTAAACAAAAGAACCCA 3'  
 30 Nor-856 5' GCCCTTGTTCTGCAGAAGAACCCA 3'

PstI

f) Arg 170 Tyr; Gly 195 Glu (R170Y;G195E (S004)):

A combination of Nor-577 and Nor-237 was performed in  
 35 analogy with the above.

g) Gly 195 Glu; 251 Glu (G195E;K251E (S018)):

A combination of Nor-237 and Nor-736 was performed in analogy with the above.

5 h) Arg 170 Tyr; Lys 251 Glu (R170Y;K251E (S011)):

A combination of Nor-577 and Nor-736 was performed in analogy with the above.

i) Arg 170 Tyr; Gly 195 Glu; Lys 251 Glu (R170Y;G195E;K251E (S012)):

A combination of Nor-577, Nor-237, and Nor-736 was performed in analogy with the above.

j) Gly 195 Glu; Lys 235 Leu (G195E;K235L):

A combination of Nor-237 and Nor-856 was performed in analogy with the above.

k) Arg 170 Tyr; Gly 195 Glu; Lys 235 Leu (R170Y;G195E;K235L):

A combination of Nor-577, Nor-237, and Nor-856 was performed in analogy with the above.

l) His 120 Asp; Lys 235 Leu (H120D;K235L (S016)):

A combination of Nor-735 and Nor-856 was performed in analogy with the above.

m) His 120 Asp; Gly 195 Glu; Lys 235 Leu (H120D;G195E;K235L (S017)):

A combination of Nor-735, Nor-237, and Nor-856 was performed in analogy with the above.

n) His 120 Asp; Arg 170 Tyr; Gly 195 Glu; Lys 235 Leu (H120D;R170Y;G195E;K235L (S019)) :

A combination of Nor-735, Nor-577, Nor-237, and Nor-856 was performed in analogy with the above.

o) His 120 Asp; Arg 170 Tyr; Gly 195 Glu; Lys 235 Leu; Lys 251 Glu (H120D;R170Y;G195E;K235L;K251E (S020)):

A combination of Nor-735, Nor-577, Nor-237, Nor-856, and Nor-736 was performed in analogy with the above.

Gapped duplex mutagenesis was performed using the plasmids pSX93, pSX119, and pSX120 as templates.

5 pSX93 is shown in Figure 3 and is pUC13 (Vieira, J. and Messing, J.: 1982, Gene 19: 259-268) harbouring an 0.7 kb XbaI-HindIII fragment of the subtilisin 309 gene including the terminator inserted in the polylinker.

For the introduction of mutations in the N-terminal  
10 part of the enzyme the plasmid pSX119 was used. pSX119 is pUC13 harbouring an EcoRI-XbaI fragment of the subtilisin 309 gene inserted into the polylinker. The templates pSX 93 and pSX119 thus cover the whole of the subtilisin 309 gene.

Plasmid pSX120 is a plasmid where the HpaI-HindIII  
15 fragment with the subtilisin 309 gene from pSX88 is inserted into EcoRV-HindIII on pDN 1681, in a way whereby the protease gene is expressed by the amy M and amy Q promoters. pDN 1681 is obtained from pDN 1380 (Diderichsen, B. and Christiansen, L.: 1988, FEMS Microbiology Letters 56: 53-60) with an inserted  
20 ted 2.85 bp ClaI fragment from B. amyloliquefaciens carrying the amy Q gene with promoter (Takkinen et al.: 1983, J. Biol. Chem. 258: 1007ff.). The construction of pSX120 is outlined in Fig. 1, showing that pDN1681 is cut with EcoR5 and HindIII, and pSX88 with HindIII and HpaI, whereafter ligation results in  
25 pSX120 regulated by the amy M and amy Q promoters.

Four further plasmids pSX170, pSX172, pSX173, and pSX186 were constructed for gapped duplex mutagenesis of the subtilisin 309 gene:

-pSX170: SphI-KpnI, 700 bp from pSX120 inserted into pUC 19  
30 SphI-KpnI, from amino acid residue 170 in mature subtilisin 309 to terminator.

-pSX172: EcoRI-SphI, 1360 bp from pSX120 inserted into pUC 19  
EcoRI-SphI, from the promoter to amino acid residue 170 in mature subtilisin 309.

35 -pSX173: like pSX170, but with G195E.

-pSX186: PvuII-EcoRI, 415 bp from pSX120 inserted into pUC 19  
HincI-EcoRI, from amino acid residue 15 to amino acid residue 206 in mature subtilisin 309.

Figure 2 shows a somewhat detailed restriction map of pSX120 on which it is indicated which fragments were used for the construction of plasmids pSX170, pSX172, pSX173, and pSX186.

- 5           The mutation a) was performed by cutting pSX93 with XbaI and ClaI as indicated in Figure 3 and described in the section "GENERATION OF SITE DIRECTED MUTATIONS IN THE SUBTILISIN GENE" and in unpublished International Patent Application no. PCT/DK 88/00002 (NOVO INDUSTRI A/S)
- 10          Mutations b), d) and e) were performed correspondingly by cutting pSX170 by SphI and KpnI.  
            Mutations f) and g) were performed as above, but with pSX173 in stead of pSX170.  
            Mutation c) was performed correspondingly by cutting  
15 pSX186 by PstI and EcoRI.  
            The mutations h) to o) were constructed by combining DNA fragments with single or double mutations b) to g) using the restriction sites NheI, XbaI, ClaI, AvaII, and KpnI as appropriate.
- 20          Further mutants were produced using similar methods or general methods as known from the literature.

#### SUBTILISIN CARLSBERG MUTANTS

- 25          For certain examples of mutations in subtilisin Carlsberg mentioned in this specification the following changes in the nucleotide sequence of the gene were introduced:

- Asp 14 Lys (D14K (C001))   (GAT -> AAG)
- Asp 120 Lys (D120K (C002))       (GAT -> AAA)
- Asp 140 Lys (D140K (C003))       (GAC -> AAA)
- 30   Asp 14 Lys + Asp 120 Lys (D14K+D120K (C004))
- Lys 27 Asp (K27D (C005)) (AAA -> GAT)
- Lys 27 Asp + Asp 120 Lys (K27D+D120K (C006))
- Asp 172 Lys (D172K (C008))       (GAC -> AAA)
- Asp 14 Lys + Asp 120 Lys + Asp 140 Lys + Asp 172 Lys  
35   (D14K+D120K+D140K+D172K (C010))
- Val 51 Asp (V51D (C100))
- Glu 54 Thr (E54T (C101))               (GGG -> ACA)
- Glu 54 Tyr (E54Y (C102))               (GGG -> TAT)

These changes were introduced by changing the corresponding oligos in the fragments concerned. The correctness of the new sequences was confirmed after which the original oligos were replaced by these new sequences and assembled into new DNA fragments. Finally the fragments were re-assembled into the new subtilisin Carlsberg gene.

#### EXPRESSION OF MUTANT SUBTILISINS

Subsequent to sequence confirmation of the correct mutation the mutated DNA fragments were inserted into plasmid pSX92 or pSX120, which were used for producing the mutants.

Plasmid pSX92 is shown in Figure 4 and was produced by cloning the Sub 309 gene into plasmid pSX62 cut at ClaI, filled in with the Klenow fragment of DNA polymerase I and cut with HindIII prior to the insertion of the fragments DraI-NheI and NheI-HindIII from the cloned Sub 309 gene.

To express the mutants the mutated fragments (XbaI-ClaI, XbaI-HindIII, or EcoRI-XbaI) were excised from the appropriate mutation plasmid pSX93, pSX119, pSX170, pSX172, pSX173, and pSX186, respectively, and inserted into pSX92, or pSX120 to obtain plasmids capable of expressing the various mutants.

The mutated pSX92 or pSX120 were then used to transform B. subtilis DN497.

The transformed cells were then spread on LB agar plates with 10 mM phosphate, pH 7, 6µg/ml chloramphenicol, and 0.2% xylose to induce the xyn-promoter in the plasmid. The plates also contained 1% skim milk so the protease producing transformants could be detected by the clear halo where the skim milk had been degraded.

After appropriate growth the mutated enzymes were recovered and purified.

#### FERMENTATION OF THE SUBTILISIN CARLSBERG SPECIES

In order to produce protease enzyme on the basis of the microorganisms carrying mutant genes for BPN' as described above, a Rushton-type Chemoferm fermenter was generally used with an eight flat blade impeller and a working volume of 8

litres. The fermenter configuration was prepared conform to the safety regulations for VMT and consisted of:

- a) A pressure controller (type 4-3122, Bell & Howell) cutting off air supply above 0.1 bar overpressure. This is done to prevent clogged exhaust air filters.
- b) A foam trap on the gas outlet made from a 20 l suction vessel having anti-foam on the bottom.
- c) A cooling water jacket without seals in order to prevent contamination of the cooling water or tap-water drain.
- d) An absolute exhaust filter is used (Gelman acro 50, 0.45 micron).
- e) Sampling via a sampling pump device with a small internal volume.

15

#### Controls

Gas flows were controlled using mass-flow meters (Brooks, type 5852, range 0-10 l).

pH was controlled using a Hartmann and Braun transmitter and a Philips controller (Witromat). Concentrated NaOH (3M) was used as a neutralizer.

Exhaust gases were analyzed using a Unor 4N (CO<sub>2</sub>) and an Oxygor 7N (O<sub>2</sub>) from Maihak, Westinghouse. Oxygen tension in the medium was determined using an industrial polarographic sterilizable oxygen probe (Ingold type 322756702).

The medium temperature was monitored using a PT100 sensor and a Honeywell temperature controller (class 84). Foaming was kept at an acceptable level using a contact electrode, while a level switch activated an anti-foam dosage pump.

All external controls were put under the control of a Hewlett Packard microcomputer (HP220).

#### Cultivation conditions

The inocula were prepared by incubating a shake flask culture at 30°C for 16h at 250 rpm in a rotary shaker (LH fermentation, type MKx). 300 ml inoculum was used for 8L medium being conditioned at the actual fermentation conditions (pH 7.0, 30°C, air flow 3.5 l/min, stirrer 1000-1500 rpm).

Dissolved oxygen concentration was kept 25% above air saturation. Anti foaming agent used was a silicon oil based material (Rhodorsil R426, Rhone Poulenc).

#### 5 Production of subtilisin protease

The (mutant) proteases were produced using the B. subtilis DB105 strain containing the mutant gene as described under gene construction. The culture medium consists of : 8 g/l NH<sub>4</sub>Cl; 4 g/l KH<sub>2</sub>PO<sub>4</sub>; 4 g/l K<sub>2</sub>HPO<sub>4</sub>; 2 g/l NaCl; 1 g/l MgSO<sub>4</sub>.2H<sub>2</sub>O; 10 g/l yeast extract; 40 g/l sucrose;; pH 7.0 and sterilized 45 min at 120°C. After sterilization 25 mg/l tryptophan; 20 mg/l Neomycin were added. Fermentations were stopped after 20 - 30 hours. The media were cleared from cells by centrifugation.

#### PROTEOLYTIC ACTIVITY OF MUTANT SUBTILISINS

The proteolytic activity of various mutants was tested against casein as protein substrate, according to the DMC method supra. The results are presented in Table IV.

From the table it is seen that mutant (S005) exhibits a slightly enhanced activity compared to the parent (S000), whereas the remaining mutants exhibit a slightly decreased activity.

TABLE IV

#### Proteolytic Activity of Mutant Subtilisins

Mutant	Relative Activity
None (S000)	100
S001	95
S003	90
S004	85
S005	105
S006	100
S012	80
S017	90
S019	70
S020	75
S024	70



WASH PERFORMANCE OF MUTANT SUBTILISINS

A:

The wash performance of various mutants in the standard liquid Detergent of pH 8.14 was tested in a model system against grass juice according to the methods detailed supra. The results are presented in table V.

Table V

		Delta R values:	
10	Mutant	Enzyme Concentration	
		1.0 mg/l	10.0 mg/l
	S000	4.0	10.7
	S001	5.9	12.8
	S003	6.0	13.5
15	S004	5.8	13.0
	S012	4.2	9.6
	S019	10.5	19.4
	S020	9.4	18.6

20 From the table it is seen that all of the tested mutants exhibited improved or equal wash performance compared to the wild type parent enzyme. The wash performance of the mutants S019 and S020 is improved so that 1.0 mg/l of these enzymes roughly stated should be able to replace 10.0 mg/l of  
25 the wild type parent enzyme, thereby indicating a substantial improvement in the wash performance for the mutant enzymes of the invention.

B:

30 The results from tests of some of the enzyme variants of the invention in the modified commercial US liquid detergent at various pH values in a model system are shown in Table VI.

72  
Table VI

Wash performance at different pH's

Mutant		Delta R					
		pI <sub>o</sub>	pH	8.0	9.0	10.0	11.0
5	S000	10.02		1.4	2.6	3.1	10.1
	S001	9.86		2.1	4.0	6.6	14.0
	S003	9.86		2.3	5.0	8.1	14.1
	S004	9.68		4.1	9.7	11.7	10.9
	S005	9.71		2.2	4.3	6.3	13.9
10	S012	9.09		5.7	11.9	13.8	6.3
	S019	9.09		6.4	10.7	12.2	3.7
	S020	6.71		7.8	10.6	8.5	2.4

The results show clearly, that shifting the pI of an enzyme in a direction where it is desired to shift the pH optimum for the wash performance of the enzyme to approach the pH of the wash liquor improves the wash performance of the enzyme.

20 D:

The wash performance of various mutants was tested against grass juice stained cotton cloths according to the method described in Assay A.

2.0 g/l of a liquid detergent (Detergent D3) was used. The detergent was dissolved in ion-exchanged water. pH was adjusted to 9.1 with NaOH/HCl.

The temperature was kept at 20°C isothermic for 10 min.

The mutants was dosed at 0.25; 0.5; 1.0; 2.0; 5.0; and 10.0 mg enzyme protein/l each.

The stability of the mutants tested was determined by measuring the denaturation temperature (maximum excess heat capacity) by differential scanning calorimetry, DSC. The heating rate was 0.5°C/min.

The stability was tested in a solution containing approx. 2 mg/ml of the mutant in 91% standard liquid detergent, the composition of which is described in Assay A. The solution was made by mixing 100 µl of enzyme solution (approx. 20 mg enzyme/ml in a buffer of 0.01 M dimethylglutaric acid, 0.002

M  $\text{CaCl}_2$ , 0.2 M  $\text{H}_3\text{BO}_3$  and 0-0.1 M NaCl pH 6.5) with 1000  $\mu\text{l}$  standard liquid detergent.

Within the group of Subtilisin 309 mutants stability results obtained by DSC are consistent with stability results 5 obtained by traditional storage stability tests.

## Results

The wash performance of various mutants in liquid 10 detergent is presented in Table VII. The results are shown as improvement factors relative to the wild type parent enzyme. The improvement factor is defined as in Assay C.

Also shown in Table VII is the denaturation tempera-  
ture in standard liquid detergent by DSC, and the difference 15 between the denaturation temperature of the wild type parent enzyme and that of the mutant in question.

Table VII

	mutant	$\text{pI}_0$	Improvement	Denaturation	Denaturation
20		calcu-	factor	temperature	temperature
		lated		by DSC	by DSC rela-
				(°C)	tive to S000
	S 000	10.06	1	65.2	0.0
	S 020	7.30	7.6	58.2	-7.0
25	S 021	9.85	1.3	69.2	+4.0
	S 022	8.07	9.3	61.9	-3.3
	S 023	8.05	8.8	63.5	-1.7
	S 024	6.86	3.9	60.6	-4.6
	S 025	8.94	6.7	69.1	+3.9
30	S 035	8.07	7.0	72.5	+7.3
	S 201	9.85	1.4	69.4	+4.2

From Table VII it is seen that all of the tested mutants exhibit improved wash performance compared to the wild 35 type parent enzyme. The best wash performance is achieved by the mutants having  $\text{pI}_0$  equal to or just below the pH of the wash solution.

Denaturation temperature by DSC shows that the stability of the single mutants S 021 (\*36D) and S 201 (N76D)

is increased by 4.0°C and 4.2°C respectively relative to the wild type parent enzyme.

Among the mutations that are incorporated in one or more of the mutants listed in Table VII it has been shown that the mutations R170Y and K251E destabilize the mutant relative to the wild type parent enzyme, whereas the mutations H120D, G195E and K235L is indifferent with respect to stability.

It is seen from Table VII that mutants containing one destabilizing mutation are destabilized, even in cases, where a stabilizing mutation is included.

The stabilizing effects of \*36D and N76D are additive. This is shown by the mutants S 025 and S 035. S 025 contains three mutations which are indifferent to stability and the stabilizing mutation \*36D. The denaturation temperature for S 025 is increased by 3.9°C relative to the wild type parent enzyme, which is equal to the increase measured for the single mutant \*36D, S 021. S 035 contains the same mutation N76D. The denaturation temperature for S 035 is increased by 7.3°C relative to the wild type parent enzyme, which, within experimental error, is equal to the sum of the increase measured for the single mutants \*36D, S 021 and N76D, S 201.

E:

The wash performance of three mutants was tested against grass juice stained cotton cloth according to the method described in Assay A.

2.0 g/l of liquid detergent D3 was used. The detergent was dissolved in ion-exchanged water. pH was adjusted to 9.1 with NaOH/HCl.

The temperature was kept at 30°C isothermic for 10 min. The mutants were dosed at 1.0 and 10.0 mg enzyme protein/l each.

## Results

The wash performance of three mutants in commercial US-liquid detergent was tested against grass juice. The results are shown in Table VIII.

Table VIII

Delta R values:

Mutant	calculated	Enzyme concentration	
	$pI_0$	1.0 mg/l	10.0 mg/l
5 S 000	10.06	4.5	13.6
S 003	9.75	9.4	18.0
S 004	9.54	13.7	18.1
S 006	9.85	6.0	15.6

10 From Table VIII it is seen that all of the mutants exhibit improved wash performance relative to the wild type parent enzyme. It is further seen that the best performance is achieved by the mutant having  $pI_0$  closest to the pH of the wash solution.

15

F:

The wash performance of two mutants was tested against grass juice stained cotton cloth according to the conditions described in Example E.

20

## Results

The wash performance of two mutants in detergent D3 was tested against grass juice stained cotton cloth. The results are shown in Table IX.

25

Table IX

Mutant	calculated	Delta R values:	
	$pI_0$	Enzyme concentration	
		1.0 mg/l	10.0 mg/l
30 S 000	10.06	5.8	15.3
S 015	9.95	8.4	20.0
S 017	9.40	17.0	20.8

35 From Table IX it is seen that all of the mutants exhibit improved wash performance relative to the wild type parent enzyme. It is further seen that the best performance is achieved by the mutant having  $pI_0$  closest to the pH of the wash solution.

G:

The wash performance of various mutants was tested on grass juice stained cotton cloth according to the method described in Assay A.

5 2.0 g/l of detergent D3 was used.

The detergent was dissolved in buffer (0,0025 M Boric acid and 0,001 M disodium hydrogen phosphate prepared in ion exchanged water). pH was adjusted to 7.0 , 8.0 , 9.0 , and 10.0 respectively with NaOH/HCl. The temperature was kept at 10 30°C isothermic for 10 min.

The mutants were dosed at 0,2 mg enzyme protein/l each.

#### Results:

15 The wash performance of some of the enzyme variants of the invention at various pH values in a model system are shown in table X.

Table X

20	Variant	Mutation	Delta R			
			pI <sub>0</sub>	pH 7.0	8.0	9.0 10.0
	S000		10.06	0.6	0.8	4.4 7.0
	S015	K235L	9.95	1.3	2.4	6.0 8.8
	S021	*36D	9.85	2.1	3.2	5.6 8.3
25	S017	H120D,G195E,K235L	9.40	2.9	5.4	10.8 14.1
	S025	*36D,H120D,R170Y, K235L	8.95	4.3	9.5	13.9 13.1
	S023	*36D,H120D,R170Y, G195E,K235L	8.05	9.6	13.0	12.4 9.2
30	S024	*36D,H120D,R170Y, G195E,K235L,K251E	6.86	9.4	10.4	6.7 4.8

The results in Table X show clearly, that shifting the pI of a protease towards the pH of the wash liquor 35 improves the wash performance of the protease.

The results also show, that all variants tested have improved performance compared to the wild type parent enzyme at pH below 10.0.

H:

The wash performance of various mutants was tested on grass juice stained cotton cloths according to the method described in Assay A.

5           2.0 g/l of liquid detergent D3 was used. The detergent was dissolved in 0,005 M glycine prepared in ion-exchanged water). pH was adjusted to 10.0, 10.25 , 10.50 , 10.75 , 11.0 , 11.5 , and 12.0, respectively, with NaOH. The temperature was kept at 30°C isothermic for 10 minutes.

10           The mutants were dosed at 0,2 mg enzyme protein/l each.

### Results:

The wash performance of some of the enzyme variants of the invention at various pH values in a model system are shown in table XI. In this case variants with slightly higher pI than the wild type parent enzyme was investigated. The pH range from pH 10.0 to 12.0 is investigated in more details than in prior examples.

20

Table XI

Variant		Mutation	Delta R				
			pI <sub>0</sub>	pH	10.0	10.25	10.50
25	S000		10.06		7.0	8.7	10.5
	S027	E89S	10.28		6.0	8.5	9.8
	S028	D181N	10.28		6.9	9.8	10.6
	S032	D197N	10.28		4.7	9.2	10.8
	S033	E271Q	10.28		7.1	6.7	7.8
	S031	D197N,E271Q	10.53		4.7	7.2	7.0

30

	Variant	Mutation	Delta R					
			pI <sub>0</sub>	pH	10.75	11.0	11.5	12.0
35	S000		10.06		12.5	14.4	10.6	3.8
	S027	E89S	10.28		11.9	14.3	12.8	5.0
	S028	D181N	10.28		13.0	14.4	10.7	4.6
	S032	D197N	10.28		13.8	13.5	11.3	5.0
	S033	E271Q	10.28		10.4	13.7	13.3	6.3
	S031	D197N,E271Q	10.53		10.7	13.0	14.4	8.7

The data in Table XI show, that at high pH values maximum performance is achieved at pH values a little above the calculated pI. Still increasing the pI of the protease tends to increase the pH of maximum performance. The effects are not as pronounced as it is seen at low pH values (assay B and G).

I:

In order to visualize the correlation between isoelectric point of the protease and the pH at which the protease has its maximum performance, the results from examples B, G, and H are used to find the pH at which each of the investigated variants (and the wild type parent enzyme) has its maximum performance. In Figure 5 this  $pH_{max}$  is shown as a function of the calculated  $pI_0$ .

Taking into account, that the pH range is investigated in steps of 1.0 pH value the correlation is obvious.

Concerning the combination of the mutants of the invention with lipase experimental results led to the following practical conclusions:

Lipase was stable for an hour in wash liquor of type O at 37°C. The presence of Savinase® led to rapid deactivation. Kazusase® led to substantially less inactivation of lipase over the period of the test.

Proteinase K was seen to be less aggressive to lipase than Savinase®, but more so than Kazusase®. Subtilisin BPN' did not however inactivate lipase at all under these conditions.

30

Preferred proteases for use e.g. in connection with lipase in wash compositions represented by type O, are mutants S001, S003, S004, S012, S019, S020, S021, S025, S035, S235.

Type O wash liquor was a 5 g/l solution at 37°C derived from the following detergent formulation (% by wt):



79

	anionic surfactant	6
	nonionic surfactant	5
	fatty acid	2.8
	acrylic polymer	3
5	zeolite	22
	carbonate	10
	sulphate	17.5
	clay	8
	tertiary amine	2
10	perborate monohydrate	13
	minors and water	to 100.

A preferred protease for use e.g. in connection with lipase in wash compositions represented by type W, is mutant  
 15 S020, S021, S025, S035, S235.

Type W wash liquor was a 2 g/l solution of a liquid detergent having the following formulation (% by wt):

20	anionic surfactant	16
	nonionic surfactant	7
	hydrotrope	6
	citric acid	6.5
	NaOH	4.1
25	monoethanolamine	2
	minors and water	to 100.

Although the present invention has been discussed and exemplified in connection with various specific embodiments  
 30 thereof this is not to be construed as a limitation to the applicability and scope of the disclosure, which extends to all combinations and subcombinations of features mentioned and described in the foregoing as well as in the attached patent claims.

PATENT CLAIMS

1. A mutated subtilisin protease, characterised in that the net electrostatic charge has been changed in comparison to the parent protease at the same pH, for example in that, in said protease there are, relative to said parent protease, fewer or more positively-charged amino acid residue(s) and/or more or fewer negatively-charged amino acid residue(s), among the amino acid residues at any one or more of positions 1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24, 25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89, 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130, 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155, 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170, 171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191, 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256, 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275, by deletion, substitution, or insertion (single or multiple) adjacent to the indicated positions, whereby said subtilisin protease has an isoelectric point ( $pI_0$ ) lower than that of said parent protease.

2. A mutated subtilisin protease in which the net electrostatic charge has been changed in comparison to the parent protease at the same pH, for example in that in said protease there are, relative to said parent protease, more or fewer positively-charged amino acid residue(s) and/or fewer or more negatively-charged amino acid residue(s), among the amino acid residues at any one or more of positions 1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24, 25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89, 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130, 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155,

156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170,  
171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191,  
192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214,  
215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242,  
5 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256,  
257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275,  
by deletion, substitution, or insertion (single or multiple)  
adjacent to the indicated positions, whereby said subtilisin  
protease has an isoelectric pH ( $pI_0$ ) higher than that of said  
10 parent protease.

3. The protease of claim 1 or 2, characterised in that  
it possesses at least one mutation affecting an amino acid  
residue occupying a position chosen from the group of posi-  
15 tions  
1, 2, 3, 4, 14, 15, 17, 18, 20, 27, 40, 41, 43, 44, 45, 46,  
51, 52, 60, 61, 62, 75, 76, 78, 79, 91, 94, 97, 100, 105, 106,  
108, 112, 113, 117, 118, 129, 130, 133, 134, 136, 137, 141,  
143, 144, 145, 146, 165, 173, 181, 183, 184, 185, 191, 192,  
20 206, 209, 210, 211, 212, 216, 239, 240, 242, 243, 244, 245,  
247, 248, 249, 251, 252, 253, 255, 256, 257, 259, 263, 269,  
271, 272,

4. The protease of claim 1, 2, or 3, characterised in  
25 that it possesses at least one mutation affecting an amino acid  
residue occupying a position chosen from the group of posi-  
tions  
1, 2, 3, 4, 14, 15, 17, 18, 20, 27, 40, 41, 43, 44, 45, 46,  
51, 52, 60, 61, 62, 75, 76, 78, 79, 91, 94, 97, 100, 105, 106,  
30 108, 112, 113, 117, 118, 129, 130, 133, 134, 136, 137, 141,  
143, 144, 145, 146, 165, 173, 181, 183, 184, 185, 191, 192,  
206, 209, 210, 211, 212, 216, 239, 240, 242, 243, 244, 245,  
247, 248, 249, 251, 252, 253, 255, 256, 257, 259, 263, 269,  
271, 272,  
35 and at least one further mutation affecting an amino acid  
residue occupying a position chosen from the group of posi-  
tions  
1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24,  
25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53,

54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89,  
 91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108,  
 109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130,  
 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155,  
 5 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170,  
 171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191,  
 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214,  
 215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242,  
 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256,  
 10 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

5. A protease as claimed in any preceding claim, further characterised in that it contains one or more of the mutations:

15 R10F, R10L, R10F+R45A+E89S+E136Q+R145A+D181N+R186P+E271Q,  
 R10F+R19Q+E89S+E136Q+R145A+D181N+E271Q+R275Q, Q12K, Q12R,  
 Q12K+P14D+T22K+N43R+Q59E+N76D+A98R+S99D+S156E+A158R+A172D+  
 N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12R+P14D+T22R+N43R+  
 Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+  
 20 S256K+S259D+A272R, Q12K+P14D+T22K+T38K+N43R+Q59E+N76D+A98R+  
 S99D+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259+  
 A272R, Q12R+P14D+T22R+T38R+N43R+Q59E+N76D+A98R+S99D+S156E+  
 A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R, Q12K+  
 P14D+T22K+T38K+N43R+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+  
 25 S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+A272R,  
 Q12R+P14D+T22R+T38R+N43R+Q59E+N76D+A98R+S99D+H120D+N140D+  
 S141R+S156E+A158R+A172D+N173K+T213R+N248D+T255E+S256K+S259D+  
 A272R, P14D, P14K, P14K+\*36D, P14K+N218D, P14K+P129D, A15K,  
 A15R, R19Q, T22K, T22R, K27R, K27V, D32\*, \*36D, \*36D+R170Y+  
 30 G195E+K251E, \*36D+H120D+R170Y+G195E+K235L, \*36D+H120D+R170Y+  
 G195E+K235L+K251E, \*36D+H120D+G195E+K235L, T38K, T38R, D41E,  
 N43R, N43K, R45A, E53R, E53K, E53G+K235L, E54G, E54Y, Q59E,  
 Q59E+N76D+A98R+S99D+S156E+A158R+A172D+N173K+T213R+N248D+  
 T255E+S256K+S259D+A272R, D60N, N76D, E89S, E89S+K251N, Y91F,  
 35 K94R, G97D, G97D+H120K, A98K, A98R, S99D, S99D+N140K, E112T,  
 H120K, H120D, H120D+K235L, H120D+G195E+ K235L, H120D+R170Y+  
 G195E+K235L, H120D+R170Y+G195E+K235L+K251E, P129D, E136Q,  
 E136K, E136R, E136Q+R10L, N140D, N140K, N140R, S141K, S141R,  
 R145A, S156E, S156E+A158R+A172D+N173K, S156E+A158R+A172D+

83

N173K+T213R, S156E+A158R+A172D+N173K+T213R+N248D+T255E+-  
 S256K+S259D+A272R, A158R, A158K, Y167V, R170Y, R170Y+G195E,  
 R170Y+K251E, R170Y+G195E+K251E, R170Y+G195E+K235L, Y171E,  
 Y171T, A172D, N173K, D181N, N184K, N184R, N185D, R186P, Y192V,  
 5 Y192V,A, G195E, G195D, G195E+T213R, G195E+K251E, G195E+K235L,  
 D197N, D197K, D197E, Q206D, Q206E, Y209L, T213R, T213K, Y214T,  
 Y214S, N218D, N218S, K235L, K235R, K237R, W241Y,L, W241Y+H249R,  
 W241L+H249R, N248D, H249R, K251R, K251E, K251N, T255E, S256R,  
 S256K, S259L, S259D, Y263W, S265K, S265R, E271Q, E271G, E271G+  
 10 K27V, E271Q,G, A272R, A272R, R275Q,  
 D14K, D14K+D120K, D14K+D120K+D140K, D14K+D120K+D140K+D172K,  
 K27D, K27D+D120K, E54T, E54Y, N97D, N97D+S98D, N97D+T213D,  
 S98D, S98D+T213D, D120K, D140K, S156E, D172K, T213D, N218D.

15 6. A protease as claimed in any preceding claim, further  
 characterised in that the mutated subtilisin protease compri-  
 ses one or more of the mutations:

	S001)	G195E
	S002)	G195D
20	S003)	R170Y
	S004)	R170Y+G195E
	S005)	K251E
	S006)	H120D
	S008)	H120D+G195E
25	S009)	T71D
	S010)	T71D+G195E
	S011)	R170Y+K251E
	S012)	R170Y+G195E+K251E
	S013)	T71D+R170Y+K251E
30	S014)	T71D+R170Y+G195E+K251E
	S015)	K235L
	S016)	H120D+K235L
	S017)	H120D+G195E+K235L
	S018)	G195E+K251E
35	S019)	H120D+R170Y+G195E+K235L
	S020)	H120D+R170Y+G195E+K235L+K251E
	S021)	*36D
	S022)	*36D+R170Y+G195E+K251E
	S023)	*36D+H120D+R170Y+G195E+K235L

	S024)	*36D+H120D+R170Y+G195E+K235L+K251E	
	S025)	*36D+H120D+G195E+K235L	
	S026)	E136R	
	S027)	E89S	
5	S028)	D181N	f
	S029)	E89S+E136R	
	S030)	E89S+D181N	0
	S031)	D197N+E271Q	
	S032)	D197N	
10	S033)	E271Q	
	S035)	*36D+N76D+H120D+G195E+K235L	
	S041)	G195F	
	S201)	N76D	
	S202)	N76D+G195E	
15	S203)	N76D+R170Y+G195E	
	S204)	H120D+G195E+K235L+K251E	
	S223)	Q59E+N76D+A98R+S99D+T213K+K235L+N248D+T255E+S256K+S259D+A272R	
	S224)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+K235L+N248D+T255E+S256K+S259D+A272R	
20	S225)	*36D+Q59E+N76D+A98R+S99D+R170Y+S156E+A158R+A172D+N173R+K235L+N248D+T255E+S256K+S259D+A272R	
	S226)	*36Q	
25	S227)	*36D+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+R170Y+G195E+K235L+N248D+T255E+S256K+S259D+A272R	
	S228)	*36D+Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R	
30	S229)	Q59E+N76D+A98R+S99D+H120D+N140D+S141R+S156E+A158R+A172D+N173K+K235L+N248D+T255E+S256K+S259D+A272R	f
	S234)	Q206D	
35	S235)	*36D+N76D	3
	S242)	*36Q+N76D+H120D+G195E+K235L	
	C001)	D14K	
	C002)	D120K	
	C003)	D140K	

	C004)	D14K+D120K
	C005)	K27D
	C006)	K27D+D120K
	C008)	D172K
5	C009)	D14K+D120K+D140K
	C010)	D14K+D120K+D140K+D172K
	C013)	N97D
	C014)	S98D
	C015)	T213D
10	C017)	S156E
	C018)	N97D+S98D
	C019)	N97D+T213D
	C022)	S98D+T213D
	C028)	N218D
15	C100)	V51D
	C101)	E54T
	C102)	E54Y

7. A mutated subtilisin protease, characterised in that  
20 it possesses an insertion mutation at position 36.

8. A protease according to claim 7, characterised in  
that it possesses an insertion mutation giving a negatively  
charged amino-acid residue at position 36, for example \*36D or  
25 \*36E.

9. A protease according to claim 7, characterised in  
that the insertion mutation provides a neutral amino-acid  
residue, e.g. a neutral polar residue, e.g. \*36A, \*36Q, or  
30 \*36N, or a positively-charged amino-acid residue, e.g. \*36R or  
\*36K.

10. A protease according to claim 7, characterised in  
that it possesses a further mutation at any one or more of  
35 positions 120, 170, 195, 235, and 251, e.g. \*36D and one or  
more of H120D, R170Y, G195E, K235L, and K251E; or \*36Q or \*36E  
and one or more of H120D, R170Y, G195E, K235L, and K251E.

11. A protease according to claim 7, characterised in that it possesses a further mutation at position 76, e.g. to substitute a negatively-charged amino acid residue at position 76, e.g. N76D or N76E.

5

12. A protease according to claim 11, characterised in that it possesses a further mutation at mutation at any one or more of positions 120, 170, 195, 235, and 251, e.g. \*36D+N76D and one or more of H120D, R170Y, G195E, K235L, and K251E; or  
10 \*36Q+N76D or \*36E+N76D and one or more of H120D, R170Y, G195E, K235L, and K251E.

13. A protease according to claim 7, characterised in that it possesses a further mutation to give a positive charge  
15 elsewhere in the protease molecule, for example at position 213, for example T213K.

14. A mutated subtilisin protease, characterized in that one or more amino acid residues in a distance of more than 15Å  
20 from the catalytic triad of said protease has been changed in comparison to the amino acid sequence of its parent enzyme, and in a way to provide for a mutant protease having an isoelectric point,  $pI_0$ , shifted in the same direction as it is desired to shift the pH optimum for the wash performance of  
25 said mutated protease.

15. A protease according to any of claims 1 to 14, characterised in having an isoelectric point ( $pI_0$ ) lower than that of the parent protease, and having one or more mutations  
30 equivalent to removal of a positively-charged amino acid residue or addition of a negatively-charged amino-acid residue, relative to the parent protease, at the site of at least one amino-acid residue located within the range of about 15Å-20Å  
35 120, or 195.

16. A protease as claimed in any preceding claim, further characterised in that it represents a mutation of a parent enzyme selected from subtilisin BPN', subtilisin amylosac-



chariticus, subtilisin 168, subtilisin mesentericopeptidase, subtilisin Carlsberg, subtilisin DY, subtilisin 309, subtilisin 147, thermitase, aqualysin, Bacillus PB92 protease, proteinase K, Protease TW7, and Protease TW3.

5

17. A protease as claimed in claim 16, further characterized in that the parent subtilisin is subtilisin 309.

18. A protease as claimed in claim 16, further characterized in that the parent subtilisin is subtilisin 147.

19. A protease as claimed in claim 16, further characterized in that the parent subtilisin is subtilisin Carlsberg.

15 20. A protease as claimed in claim 16, further characterized in that the parent subtilisin is Bacillus PB92 protease.

21. A method for determining or selecting the position(s) and the amino acid(s) to be deleted, substituted or  
20 inserted for the amino acid(s) in a parent enzyme, characterized in that the selection is performed in a way whereby the calculated net electrostatic charge (=NEC) in a contemplated mutant enzyme has been changed in comparison to the NEC in the parent enzyme of choice calculated at the same pH value.

25

22. The method of claim 21, further characterized in that the amino acid residues selected are situated on, at or near to the surface of said parent enzyme.

30 23. The method of claim 21 or 22, further characterized in that the amino acid residues selected are chosen among the amino acid residues at any one or more of positions

1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24,  
25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53,  
35 54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89,  
91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108,  
109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130,  
131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155,  
156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170,

171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191,  
192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214,  
215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242,  
243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256,  
5 257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

24. The method of claim 21, 22 or 23, further characterized in that the amino acid residues selected are chosen among the amino acid residues at any one or more of positions  
10 1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24,  
25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53,  
54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89,  
91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108,  
109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130,  
15 131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155,  
156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170,  
171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191,  
192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214,  
215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242,  
20 243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256,  
257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

25. The method of claim 21, 22, 23 or 24, further characterized in that the amino acid residues selected are chosen to  
25 affect an amino acid residue occupying a position chosen from the group of positions

1, 2, 3, 4, 14, 15, 17, 18, 20, 27, 40, 41, 43, 44, 45, 46,  
51, 52, 60, 61, 62, 75, 76, 78, 79, 91, 94, 97, 100, 105, 106,  
108, 112, 113, 117, 118, 129, 130, 133, 134, 136, 137, 141,  
30 143, 144, 145, 146, 165, 173, 181, 183, 184, 185, 191, 192,  
206, 209, 210, 211, 212, 216, 239, 240, 242, 243, 244, 245,  
247, 248, 249, 251, 252, 253, 255, 256, 257, 259, 263, 269,  
271, 272,

and at least one further affecting an amino acid residue  
35 occupying a position chosen from the group of positions

1, 2, 3, 4, 6, 9, 10, 12, 14, 15, 17, 18, 19, 20, 21, 22, 24,  
25, 27, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 50, 51, 52, 53,  
54, 55, 56, 57, 58, 59, 60, 61, 62, 75, 76, 77, 78, 79, 87, 89,  
91, 94, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108,

109, 112, 113, 115, 116, 117, 118, 120, 126, 128, 129, 130,  
131, 133, 134, 136, 137, 140, 141, 143, 144, 145, 146, 155,  
156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170,  
171, 172, 173, 181, 182, 183, 184, 185, 186, 188, 189, 191,  
5 192, 194, 195, 197, 204, 206, 209, 210, 211, 212, 213, 214,  
215, 216, 217, 218, 235, 236, 237, 238, 239, 240, 241, 242,  
243, 244, 245, 247, 248, 249, 251, 252, 253, 254, 255, 256,  
257, 259, 260, 261, 262, 263, 265, 269, 271, 272, 275.

10 26. A method as claimed in any of the claims 21 to 25,  
further characterised in that the parent enzyme is selected  
from subtilisin BPN', subtilisin amylosacchariticus, subtili-  
sin 168, subtilisin mesentericopeptidase, subtilisin Carls-  
berg, subtilisin DY, subtilisin 309, subtilisin 147, thermi-  
15 tase, Bacillus PB92 protease, proteinase K, protease TW7, and  
protease TW3.

27. A method as claimed in claim 26, further characteri-  
sed in that the parent subtilisin is subtilisin 309.

20

28. A method as claimed in claim 26, further characteri-  
sed in that the parent subtilisin is subtilisin 147.

29. A method as claimed in claim 26, further characteri-  
25 sed in that the parent subtilisin is subtilisin Carlsberg.

30. A method as claimed in claim 26, further characteri-  
sed in that the parent subtilisin is Bacillus PB92 protease.

1/5




Fig. 1

2/5





Fig. 2

3/5

**Fig. 3**

4/5




Fig. 4

5/5





Fig. 5



# INTERNATIONAL SEARCH REPORT

International Application No. PCT/DK 90/00164

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC5: C 12 N 9/50, C 12 N 15/57, C 11 D 7/42		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched		
Classification System	Classification Symbols	
IPC5	C 12 N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched <sup>8</sup>		
SE,DK,FI,NO classes as above		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
P,X	WO, A1, 8906279 (NOVO INDUSTRI A/S) 13 July 1989, see the claims --	1-20
P,X	US, A, 4760025 (DAVID A. ESTELL ET AL.) 26 July 1988, see example 15 and 16 --	1-10
P,X	WO, A1, 8909819 (GENEX CORPORATION) 19 October 1989, see the claims --	1-10
P,X	WO, A1, 8909830 (GENEX CORPORATION) 19 October 1989, see page 10 --	1-20
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><b>* Special categories of cited documents:</b> <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
2nd October 1990	1990-10-09	
International Searching Authority	Signature of Authorized Officer	
SWEDISH PATENT OFFICE	 Yvonne Siösteen	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X	WO, A1, 8808033 (AMGEN INC.) 20 October 1988, see page 9, line 10-21 --	1-30
X	J. Mol. Biol., Vol. 193, 1987 Alan J Russel et al.: "Electrostatic Effects on Modification of Charged Groups in the Active Site Cleft of Subtilisin by Protein Engineering ", see page 803 - page 813 see table 5 --	1-30
X	NATURE, Vol. 318, November 1985 Paul G. Thomas et al.: "Tailoring the pH dependence of enzyme catalysis using protein engineering ", see page 375 - page 376 and the whole document --	1,2,23, 24
X	Chemical Abstracts, volume 106, no. 21, 25 May 1987, (Columbus, Ohio, US), Wells, J.A. et al.: "Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin ", see page 333, abstract 171899m, & Philos. Trans. R. Soc. London, A 1986, 317(15), 415- 423 --	1-30
X	NATURE, Vol. 330, November 1987 Michael J.E. Sternberg et al.: "Prediction of electrostatic effects of engineering of protein charges ", see page 86 - page 88 --	1,2,23, 24
X	NATURE, Vol. 328, August 1987 Alan J. Russell et al.: "Rational modification of enzyme catalysis by engineering surface charge ", see page 496 - page 500 -- -----	1,2,23, 24

ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.PCT/DK 90/00164

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 90-08-28. The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 8906279	89-07-13	NONE	
US-A- 4760025	88-07-26	AU-B- 587960	89-09-07
		AU-D- 2956884	85-01-03
		AU-D- 3720889	89-12-07
		EP-A- 0130756	85-01-09
		EP-A- 0246678	87-11-25
		EP-A- 0247647	87-12-02
		EP-A- 0357157	90-03-07
WO-A1- 8909819	89-10-19	NONE	
WO-A1- 8909830	89-10-19	NONE	
WO-A1- 8808033	88-10-20	AU-D- 1701688	88-11-04
		EP-A- 0309565	89-04-05
		US-A- 4914031	90-04-03